

# On star and caterpillar arboricity

Daniel Gonçalves<sup>a</sup>, Pascal Ochem<sup>b</sup>

<sup>a</sup>*LIRMM UMR 5506, CNRS, Université Montpellier 2,  
161 rue Ada, 34392 Montpellier Cedex 5, France*

<sup>b</sup>*LRI UMR 8623, CNRS, Université Paris-Sud,  
Bât 490, 91405 Orsay Cedex, France*

---

## Abstract

We give new bounds on the star arboricity and the caterpillar arboricity of planar graphs with given girth. One of them answers an open problem of Gyárfás and West: there exist planar graphs with track number 4. We also provide new NP-complete problems.

*Key words:* NP-completeness, partitioning problems, edge coloring

---

## 1 Introduction

Many graph parameters in the literature are defined as the minimum size of a partition of the edges of the graph such that each part induces a graph of a given class  $C$ . The most common is the chromatic index  $\chi'(G)$ , in this case  $C$  is the class of graphs with maximum degree one. Vizing [18] proved that  $\chi'(G)$  either equals  $\Delta(G)$  or  $\Delta(G) + 1$ , where  $\Delta(G)$  denotes the maximum degree of  $G$ . Deciding whether  $\chi'(G) = 3$  is shown to be NP-complete for general graphs in [13]. The arboricity  $a(G)$  is another well studied parameter, for which  $C$  is the class of forests. In [15], Nash-Williams proved that:

$$a(G) = \max_{H \subseteq G} \left\lceil \frac{|E(H)|}{|V(H)| - 1} \right\rceil \quad (1)$$

with the maximum being over all the subgraphs  $H = (E(H), V(H))$  of  $G$ . Even with this nice formula, the polynomial algorithm computing the arboricity of

---

*URLs:* [goncalves@lirmm.fr](mailto:goncalves@lirmm.fr) (Daniel Gonçalves), [ochem@lri.fr](mailto:ochem@lri.fr) (Pascal Ochem).

a graph is not trivial [12]. Other similar parameters have been studied. A *star* is a tree of diameter at most two. A *caterpillar* is a tree whose non-leaf vertices form a path. For the *star arboricity*  $sa(G)$  and the *caterpillar arboricity*  $ca(G)$ , the corresponding class  $C$  is respectively the class of star forests and the class of caterpillar forests. Since stars are caterpillars which are trees, and since trees are easily partitionable into two star forests, we have the following two inequalities for any graph  $G$ .

$$sa(G) \geq ca(G) \geq a(G) \tag{2}$$

$$2a(G) \geq sa(G) \tag{3}$$

A proper vertex coloring of a graph is *acyclic* if there is no bicolored cycle. Let  $\chi_a(G)$  denote the acyclic chromatic number of a graph  $G$ . Hakimi *et al.* [11] showed the following relation between  $\chi_a(G)$  and  $sa(G)$ .

**Theorem 1 (Hakimi *et al.*)** *For any graph  $G$ , we have  $\chi_a(G) \geq sa(G)$ .*

Other interesting graph parameters include the track number  $t(G)$  [10,14] and the subchromatic index  $\chi'_{sub}(G)$  [6]. For the track number,  $C$  is the class of interval graphs. For the subchromatic index,  $C$  is the class of graphs whose connected components are stars or triangles. Notice that the class of triangle-free interval graphs is equivalent to the class of caterpillar forests. Thus, if  $G$  is triangle-free, then  $t(G) = ca(G)$  and  $\chi'_{sub}(G) = sa(G)$ .

Given a tree  $T$ , a  $T$ -free forest is a forest without subgraphs isomorphic to  $T$ . For example, the  $P_n$ -free forests and the  $K_{1,n}$ -free forests correspond to, respectively, the forests with diameter at most  $n - 2$  and to the forests with degree at most  $n - 1$ . Given a tree  $T$ , the  $T$ -free arboricity  $T$ - $fa(G)$  of a graph  $G$  is the minimum number of  $T$ -free forests needed to cover the edges of  $G$ . In this case  $C$  is the class of  $T$ -free forests. Using this terminology, we can redefine some of the parameters we introduced. For  $n \geq 2$ , let  $S_n$  be the tree arised from  $K_{1,n}$  by subdividing each edge once. The chromatic index, the star arboricity, and the caterpillar arboricity correspond to, respectively, the  $P_3$ -free arboricity, the  $P_4$ -free arboricity, and the  $S_3$ -free arboricity.

If a tree  $T_1$  is a subtree of a tree  $T_2$ , then  $T_1$ - $fa(G) \geq T_2$ - $fa(G)$ . So, the poset of trees produces a poset of arboricities. For example, since  $P_4 \subset S_2 \subset S_3 \subset \dots \subset S_n$ , we have  $P_4$ - $fa(G) \geq S_2$ - $fa(G) \geq S_3$ - $fa(G) \geq \dots \geq S_n$ - $fa(G)$ , for any graph  $G$ .

In [11], it is proved that deciding whether a graph  $G$ , satisfies  $sa(G) \leq 2$  is NP-complete. We obtain the same result for a very restricted graph class.

**Theorem 2** *For any  $g \geq 3$ , deciding whether a bipartite planar graph  $G$  with girth at least  $g$  and maximum degree 3 satisfies  $sa(G) \leq 2$  is NP-complete.*

This implies that there exist planar graphs of arbitrarily large girth with star arboricity at least 3. This lower bound is tight for  $g \geq 7$ . Since planar graphs of girth  $g \geq 7$  are acyclically 3-colorable [2], their star arboricity is at most 3 by Theorem 1.

As we already mentioned, if  $G$  is triangle-free, then  $sa(G) = \chi'_{sub}(G)$ . So, this theorem answers a question of Fiala and Le [6].

**Corollary 3** *Deciding whether a planar graph  $G$  satisfies  $\chi'_{sub}(G) \leq 2$  is NP-complete.*

Let us denote by  $L(G)$  the line graph of  $G$  and by  $\mathcal{L}$  the class of line graphs of “planar bipartite graphs with maximum degree three and girth at least six”. Notice that graphs in  $\mathcal{L}$  are planar with maximum degree four and line graphs of bipartite graphs, thus perfect [3]. This class of graph is very restricted, it corresponds to planar  $(K_{1,3}, K_4, K_4^-, C_4, \text{odd-hole})$ -free graphs. The complexity of determining the subchromatic number of a graph is an interesting question. Deciding whether a graph  $G$  satisfies  $\chi_{sub}(G) \leq 2$  is NP-complete if  $G$  is planar [8] or if  $G$  is perfect [4]. Theorem 2 shows that it is also the case for perfect planar graphs since  $\chi'_{sub}(G) = \chi_{sub}(L(G))$ .

**Corollary 4** *Deciding whether a graph  $G \in \mathcal{L}$  satisfies  $\chi_{sub}(G) \leq 2$  is NP-complete.*

A graph is *2-degenerate* if all of its subgraphs contain a vertex of degree at most 2.

**Theorem 5** *Deciding whether a 2-degenerate bipartite planar graph  $G$  satisfies  $sa(G) \leq 3$  is NP-complete.*

Shermer [17] proved that it is NP-complete to decide whether a graph  $G$  has caterpillar arboricity 2. We generalize here his result to  $S_n$ -free arboricity and consider more restricted graph classes.

**Theorem 6** *The following problems are NP-complete :*

- (1) *For every  $n \geq 2$ , deciding whether a 2-degenerate bipartite planar graph  $G$  satisfies  $S_n$ -fa( $G$ )  $\leq 3$ .*
- (2) *For every  $n \geq 3$ , deciding whether a 2-degenerate bipartite planar graph  $G$  of girth  $g \geq 6$  satisfies  $S_n$ -fa( $G$ )  $\leq 2$ .*

Theorem 6.1 implies the existence of bipartite planar graphs with caterpillar arboricity four and, as we already mentioned, the track number of a triangle-free graph equals its caterpillar arboricity. It is proved in [10] that deciding whether a graph  $G$  has track number  $t(G) \leq k$  is NP-complete for  $k = 2$  and conjectured that it is also the case for higher  $k$ . Here we proved that it is the

case for a restricted family of graphs and for  $k = 2$  or  $3$ .

**Corollary 7** *It is NP-complete to decide whether a 2-degenerate bipartite planar graph  $G$  satisfy  $t(G) \leq 2$  (resp.  $t(G) \leq 3$ ).*

The *interval number*  $i(G)$  is the smallest  $k$  such that every vertex of the graph  $G$  can be represented as a set of at most  $k$  intervals of a line and there is an edge  $uv$  iff the segments of  $u$  and  $v$  intersect. Scheinerman and West [16] proved that the interval number of planar graphs is at most 3 and the first author [9] proved that the caterpillar arboricity of planar graphs is at most 4. This implies that the maximum track number of planar graphs is either 3 or 4. In [14], Kostochka and West proved that the maximum track number of outerplanar graphs equals their maximum interval number, 2. We can deduce from Theorem 6.1 that this is not the case for planar graphs, which answers an open question of Gyárfás and West [10].

**Corollary 8** *There exist bipartite planar graphs with track number four.*

Theorem 6.2 implies that there exist planar graphs of girth  $g \geq 6$  with caterpillar arboricity at least three. The next theorem shows that this lower bound is tight.

**Theorem 9** *For any planar graph  $G$  with girth  $g \geq 6$ ,  $ca(G) \leq 3$ .*

Contrarily to the star arboricity, the caterpillar arboricity of planar graphs with sufficiently large girth is two.

**Theorem 10** *For any planar graph  $G$  with girth  $g \geq 10$ ,  $ca(G) \leq 2$ .*

In the next section, we define the notion of  $T$ -fa coloring, which is needed in the proofs. Section 3 and 4 are devoted to the proofs of the upper bounds and of the NP-completeness results, respectively.

## 2 $T$ -free arboricity and $T$ -fa coloring

We define the  $T$ -fa colorings for  $T = P_4$  and  $T = S_n$  with  $n \geq 2$ .

**Definition 11** *For  $T = P_4$  (resp.  $T = S_n$  with  $n \geq 2$ ), a  $k$ - $T$ -fa coloring of a graph  $G$  is a  $k$ -edge-coloring of  $G$  and a partial orientation of its edges such that:*

- *The graph induced by a color class is a  $T$ -free forest.*
- *If the edge  $uv$  is colored  $i$  and is oriented towards  $v$ , then  $v$  is a leaf in the  $i^{\text{th}}$  forest  $F_i$  (the  $T$ -free forest induced by the edges colored  $i$ ).*

- The graph induced by the unoriented edges has maximum degree 0 (resp.  $n - 1$ ).

It is clear that if a graph  $G$  has a  $k$ - $T$ - $fa$  coloring, then  $T$ - $fa(G) \leq k$  (by the first point of the definition). The reverse also holds. Indeed, given an edge partition of  $G$  into  $k$   $T$ -free forests we can construct a  $k$ - $T$ - $fa$  coloring of  $G$ . For this, color each edge set with a given color and then orient to a leaf of  $F_i$  each edge of  $F_i$  incident to leaf. With this construction, any vertex  $v$  incident to an unoriented edge colored  $i$  is incident to another edge colored  $i$ . So, if there was a vertex incident to an (resp.  $n - 1$ ) unoriented edge(s) colored  $i$ , this would contradict the “ $T$ -freeness” of the partition.

If the graph  $G$  is  $k$ - $T$ - $fa$  colored, for each of its  $k$  forests, we distinguish two types of vertices. The *ends*, which have an incident arc in this forest oriented toward them, and the *inner vertices*. A  $k$ - $T$ - $fa$  coloring of  $G$  is *suitable* if every vertex of  $G$  is an end in at most one forest (*i.e.* is an inner vertex in  $k - 1$  or  $k$  forests).

### 3 Upper bounds on the caterpillar arboricity

The maximum average degree  $mad(G)$  of a graph  $G$  is defined by  $mad(G) = \max\{2|E(H)|/|V(H)|, H \subseteq G\}$ . Since we consider planar graphs, we will use the following well known observation based on Euler’s formula:

**Lemma 12** *If  $G$  is a planar graph with girth at least  $g$ , then  $mad(G) < \frac{2g}{g-2}$ .*

Theorems 9 and 10 will be deduced, using this lemma, from a proposition of the form “every graph  $G$  of girth at least  $g$  with  $mad(G) < q = \frac{2g}{g-2}$  has a suitable  $k$ - $P$ - $fa$  coloring”. The proof of these propositions is based on the *discharging method*, as used in [1]. We consider a graph  $H$  of girth at least  $g$  that has no suitable coloring and is minimal for the subgraph partial order. This means that every proper subgraph  $H'$  of  $H$  has a suitable coloring.

First, we provide a set  $S$  of configurations that  $H$  cannot contain due to its minimality property. To show that a configuration  $C \in S$  is forbidden, we suppose that  $H$  contains  $C$  and then argue that any suitable coloring of some proper subgraph of  $H$  can be extended in a suitable coloring of the whole graph  $H$ , which is a contradiction. Then we have to prove that any graph  $K$  avoiding every configuration in  $S$  satisfies  $mad(K) \geq q$ . We assume that every vertex  $v$  is assigned an initial charge equal to its degree  $d(v)$  and define a *discharging procedure* that preserves the total charge of the graph. We show that if the discharging procedure is applied to a graph  $K$  avoiding  $S$ , then the

final charge  $d^*(v)$  of every vertex  $v \in V(K)$  satisfies  $d^*(v) \geq q$ . We thus have

$$mad(K) \geq \frac{2|E(K)|}{|V(K)|} = \frac{\sum_{v \in V(K)} d(v)}{|V(K)|} = \frac{\sum_{v \in V(K)} d^*(v)}{|V(K)|} \geq \frac{q|V(K)|}{|V(K)|} = q.$$

In every figure depicting forbidden configurations, every neighbor of a “white” vertex is drawn, whereas a “black” vertex may have other neighbors in the graph. Two or more black vertices may coincide in a single vertex, provided they do not share a common white neighbor.

It is for example clear that for  $k \geq 2$ , if  $H$  is a minimal graph having no suitable  $k$ - $S_3$ - $fa$  coloring, then its minimum degree is at least 2. If there was a 1-vertex  $v$  we could easily extend any suitable  $k$ - $S_3$ - $fa$  coloring of  $H \setminus \{v\}$ .

### 3.1 Proof of Theorem 9

**Lemma 13** *Let  $H$  be a minimal graph of girth at least 6 having no suitable 3- $S_3$ - $fa$  coloring. Then  $H$  does not contain a 2-vertex adjacent to a  $\leq 5$ -vertex.*

**Proof.** Suppose that  $H$  contains a 2-vertex  $u$  adjacent to both a  $\leq 5$ -vertex  $v$  and a vertex  $w$ . Consider a suitable coloring of  $H \setminus \{u\}$  into three forests  $F_1, F_2, F_3$ . In this coloring, the vertex  $v$  is an inner vertex in at least two forests and has degree at most 4 in  $H \setminus \{u\}$ . This implies that there is a forest, say  $F_1$ , in which  $v$  is an inner vertex and such that  $v$  is incident to at most 1 unoriented edges of  $F_1$ . The vertex  $w$  is an inner vertex in at least two forests. Let  $F_i$  be one of these forests with  $i \neq 1$ . Now we can extend the coloring to  $H$  by coloring the edges  $uv$  and  $uw$  respectively 1 and  $i$ , letting  $uv$  unoriented, and orienting  $uw$  towards  $u$ . This  $S_3$ - $fa$  coloring is suitable since  $u$  is just an end in  $F_i$ .  $\square$

We apply the following discharging rule to the graph  $H$  considered in Lemma 13: each  $\geq 6$ -vertex gives  $\frac{1}{2}$  to each of its 2-neighbors. Let us check that for every  $v \in V(H)$ ,  $d^*(v) \geq 3$ :

- $d(v) = 2$ :  $v$  has two  $\geq 6$ -neighbors by Lemma 13, so  $d^*(v) = 2 + 2 \cdot \frac{1}{2} = 3$ .
- $d(v) = k$ ,  $3 \leq k \leq 5$ : the charge of  $v$  is unchanged, so  $d(v) = d^*(v) = k \geq 3$ .
- $d(v) = k \geq 6$ :  $d^*(v) \geq k - k \cdot \frac{1}{2} = \frac{k}{2} \geq 3$ .

This shows that the maximum average degree of a minimal graph of girth at least 6 having no suitable 3- $S_3$ - $fa$  coloring is at least 3. By Lemma 12, we thus have that every planar with girth at least 6 has a suitable 3- $S_3$ - $fa$  coloring, which proves Theorem 9

### 3.2 Proof of Theorem 10

**Lemma 14** *Let  $H$  be a minimal graph of girth at least 10 having no suitable  $2$ - $S_3$ -fa coloring. Then  $H$  does not contain any of the configurations depicted in Figure 1.*




Fig. 1. Forbidden configurations for Lemma 1.

**Proof.** We consider each of the configurations:

- (i) Suppose  $H$  contains the configuration (i) depicted in Figure 1. Consider a suitable  $2$ - $S_3$ -fa coloring of  $H \setminus \{y\}$ . In every case,  $z$  is an inner vertex in some forest  $F_i$  such that  $z$  is incident to at most one non-oriented edge colored  $i$ . We can extend this coloring to  $H$  such that  $xy$  and  $yz$  are non-oriented,  $vx$  and  $xy$  get different colors, and  $yz$  gets color  $i$ .
- (ii) Suppose  $H$  contains the configuration (ii) depicted in Figure 1. Consider a suitable  $2$ - $S_3$ -fa coloring of the graph  $H'$  obtained from  $H$  by deleting the edge  $yz$ . We can always modify this coloring into a suitable  $2$ - $S_3$ -fa coloring of  $H'$  such that  $xy$  is non-oriented and there exist no monochromatic path connecting  $y$  to any  $u_i$ . In every case,  $z$  is an inner vertex in some forest  $F_i$  such that  $z$  is incident to at most one non-oriented edge colored  $i$ . We can extend this coloring to  $H$  such that  $yz$  is non-oriented and gets color  $i$ .

□

A 3-vertex is *weak* if it has three 2-neighbors. A 2-vertex is *weak* if is adjacent to a 2-vertex or a weak 3-vertex. We apply the following discharging rules to the graph  $H$  considered in Lemma 14: each  $\geq 4$ -vertex gives  $\frac{1}{2}$  to its weak 2-neighbors and  $\frac{1}{4}$  to its non-weak 2-neighbors, each non-weak 3-vertex gives  $\frac{1}{4}$  to its 2-neighbors. Let us check that for every  $v \in V(H)$ ,  $d^*(v) \geq \frac{5}{2}$ :

- $d(v) = 2$ : if  $v$  is weak, then  $v$  has a  $\geq 4$ -neighbor (see Figure 1.(i) and Figure 1.(ii) with  $m = 2$ ), so  $d^*(v) = 2 + \frac{1}{2} = \frac{5}{2}$ . Otherwise  $v$  receives  $\frac{1}{4}$  from each neighbor, so  $d^*(v) \geq 2 + 2 \cdot \frac{1}{4} = \frac{5}{2}$ .
- $d(v) = 3$ : if  $v$  is weak, then  $d^*(v) = d(v) = 3 > \frac{5}{2}$ . Otherwise  $v$  has at most two 2-neighbors, so  $d^*(v) \geq 3 - 2 \cdot \frac{1}{4} = \frac{5}{2}$ .
- $d(v) = 4$ : if  $v$  has four 2-neighbors, then its 2-neighbors are not weak (see Figure 1.(ii) with  $m = 3$ ), so  $d^*(v) \geq 4 - 4 \cdot \frac{1}{4} = 3 > \frac{5}{2}$ . Otherwise  $v$  has at most three 2-neighbors, so  $d^*(v) \geq 4 - 3 \cdot \frac{1}{2} = \frac{5}{2}$ .

-  $d(v) = k \geq 5: d^*(v) \geq k - k\frac{1}{2} = \frac{k}{2} \geq \frac{5}{2}$ .

This shows that the maximum average degree of a minimal graph of girth at least 10 having no suitable  $2-S_3$ -fa coloring is at least  $\frac{5}{2}$ . By Lemma 12, we thus have that every planar with girth at least 10 has a suitable  $2-S_3$ -fa coloring, which proves Theorem 10

#### 4 NP-completeness results

Theorems 5 and 6.1 are each obtained by a polynomial reduction from the problem 3-COLORABILITY which is NP-complete on planar graphs with maximum degree 4 [7]. A *subcoloring* of a graph is a partition of its vertex set such that each part induces a disjoint union of cliques. Theorems 2 and 6.2 are each obtained by a polynomial reduction from the problem 2-SUBCOLORABILITY which is NP-complete on triangle-free planar graphs with maximum degree 4 [5,8]. Notice that on triangle-free graphs, a 2-subcoloring corresponds to a vertex partition into two graphs with maximum degree 1. Let us now describe the reductions for Theorems 5 and 6.1 (resp. Theorems 2 and 6.2). Given a planar graph (resp. a triangle-free planar graph)  $G$ , we construct a graph  $G'$  that belong to the class specified in the theorem as follows: we add a “vertex gadget” to every vertex  $v$  of  $G$  and replace every edge  $uv$  of  $G$  by an “edge gadget”. The vertex gadget forces the  $v$  to be an inner vertex in at most one forest  $F_i$  for any  $k$ - $T$ -fa coloring (with  $k$  and  $T$  as mentioned in the theorem). The edge gadget is such that  $G'$  is  $k$ - $T$ -fa colorable if and only if  $G$  is 3-colorable (resp. 2-subcolorable). More precisely if  $G$  has a vertex coloring  $c$ , then  $G'$  has  $k$ - $T$ -fa coloring such that every original vertex  $v$  is an inner vertex in  $F_{c(v)}$ , and conversely, if  $G'$  has a  $k$ - $T$ -fa coloring such that every original vertex  $v$  is an inner vertex of  $F_i$ , then taking  $c(v) = i$  gives a vertex coloring of  $G$ .

##### 4.1 Proof of Theorem 2




Fig. 2. The vertex gadget and its 2- $P_4$ -fa colorings.



Let  $C'_n$  (resp.  $P'_n$ ) be the graph obtained from the cycle  $C_n$  (resp. the path  $P_n$ ) by adding, for each vertex  $v \in V(C_n)$  (resp.  $v \in V(P_n)$ ), a new vertex  $v'$  and an edge  $vv'$ . Note that in a  $2-P_4$ -fa coloring, if a vertex  $v$  of degree at least three has an incident edge oriented to  $v$  and colored 1, then all the remaining edges incident to  $v$  are colored 2 and are oriented from  $v$  to the other end. This implies that in any  $2-P_4$ -fa coloring of  $C'_{2n}$  each vertex of degree one is incident to an edge oriented toward it. The vertex gadget is the graph depicted in Figure 2 obtained from  $C'_{2n}$  and  $P'_{18n-1}$ . This graph is bipartite and the size of the vertices in the figure indicate in which set of the bipartition they are. In any  $2-P_4$ -fa coloring of the vertex gadget, the edge  $xy$  is oriented toward  $y$ . This imply that in any of these  $2-P_4$ -fa colorings there is a  $i \in \{1, 2\}$  such that at most one of the vertices  $z_{2n}, z_{4n}, z_{6n}, \dots, z_{16n}$  ( $z_4$  in Figure 2) is not an end of  $F_i$ . Furthermore, for any  $j$  with  $1 \leq j \leq 8$ , there is a  $2-P_4$ -fa coloring in which the vertex  $z_{2jn}$  is an end of  $F_{3-i}$ . There are also  $2-P_4$ -fa colorings of the vertex gadget in which all the vertices  $z_{2jn}$ , for  $1 \leq j \leq 8$ , are ends of  $F_i$ .




Fig. 3. The connection between two vertex gadgets.

Given a triangle-free planar graph  $G$  with maximum degree 4, we construct  $G'$  by replacing every vertex  $v$  of  $G$  by a copy of the vertex gadget, denoted  $H_v$ . Every vertex  $v$  of  $G$  numbers its incident edges from 1 to  $\deg(v) \leq 4$  going around  $v$  in the clockwise sense. For every edge  $uv$  of  $G$  we connect  $H_u$  and  $H_v$  in the following way. Let  $i_u$  (resp.  $i_v$ ) be the number of  $uv$  with respect to  $u$  (resp.  $v$ ). Identify the vertices  $z_{(2i_u-1)2n}$  and  $z_{(2i_u)2n}$  of  $H_u$  respectively with the vertices  $z_{(2i_v)2n}$  and  $z_{(2i_v-1)2n}$  of  $H_v$ . In Figure 3.(a) we have  $i_u = 3$  and  $i_v = 1$  and the connection of  $H_u$  and  $H_v$  is depicted in Figure 3.(b) or (c). The graph  $G'$  is planar, bipartite, with maximum degree three and may have arbitrary girth (its girth is  $2n$ ). We now have to show that  $\chi_{sub}(G) \leq 2$  if and only if  $sa(G') \leq 2$ .

Given a 2-subcoloring  $c$  of  $G$  we obtain a  $2-P_4$ -fa coloring of  $G'$  by coloring each vertex gadget  $H_u$  in such way that most of the vertices  $z_{2in}$  are ends in  $F_{c(u)}$ . If  $u$  has no neighbor  $v$  such that  $c(u) = c(v)$ , then all its vertices  $z_{2in}$  are ends in  $F_{c(u)}$ . If  $u$  has a neighbor  $v$  such that  $c(u) = c(v)$ , let  $i_u$  be the number of the edge  $uv$  with respect to  $u$ . By definition of a 2-subcoloring  $u$  has at most one such neighbor. In this case let the vertex  $z_{(2i_u)2n}$  be an end in  $F_{3-c(u)}$ . This coloring of  $G'$  is a  $2-P_4$ -fa coloring. Indeed, if an edge  $uv$  of  $G$  is

such that  $c(u) \neq c(v)$ , then we see in Figure 3.(b) that the  $2-P_4$ -fa colorings of  $H_u$  and  $H_v$  fit. If an edge  $uv$  of  $G$  is such that  $c(u) = c(v)$ , then we see in Figure 3.(c) that the  $2-P_4$ -fa colorings of  $H_u$  and  $H_v$  also fit.

Conversely, given a  $2-P_4$ -fa coloring of  $G'$  we obtain a 2-subcoloring of  $G$  by coloring each vertex  $u$  of  $G$  with the color  $i \in \{1, 2\}$  that verify: most of the vertices  $z_{2jn}$  of  $H_u$  are ends of  $F_i$ . Since there is at most one vertex  $z_{2jn}$  that is not an end of  $F_i$ , the vertex  $u$  has at most one neighbor in  $G$  with the same color.

#### 4.2 Proof of Theorem 5




Fig. 4. The vertex gadget and the edge gadget for the reduction of Theorem 5.

Given a planar graph  $G$ , we construct  $G'$  by adding to every vertex of  $G$  the vertex gadget depicted in Figure 4 and by subdividing every edge of  $G$ . The graph  $G'$  is clearly planar, bipartite and 2-degenerated.




Fig. 5.  $3-P_4$ -fa coloring of  $\mathcal{A}$  with conditions on  $x_1$  and  $x_2$ .

First, we comment on how to  $3-P_4$ -fa color the graph  $\mathcal{A}$  depicted in Figure 5. Notice that if a vertex has two incoming edges colored 1 and 2, all its remaining incident edges have to be colored 3. In the first drawing, we impose that all the edges  $x_i y_j$  are oriented toward  $y_j$ . This implies that all the edges  $y_i z_j$  are oriented toward  $z_j$ , and that we just used two colors for these edges. This finally implies that all the remaining edges incident to the  $z_i$ 's have the same color, which is not allowed since each color induces a forest. In the second drawing, we impose that just one edge  $x_i y_j$  is oriented toward  $x_2$  and that the edges incident to  $x_1$  have the same color, 1. The edges  $x_2 y_1$  and  $y_2 x_2$  have to be respectively colored 2 and 3. This implies that the edges  $y_1 z_i$  are oriented toward  $z_i$  and colored 3. This implies that the edges  $y_2 z_i$  are oriented toward  $z_i$  and colored 2. This finally implies that all the remaining edges incident to the  $z_i$ 's have the same color, which is not allowed. In the third drawing, we

impose that just one edge  $x_i y_j$  is oriented toward  $x_2$ , that the edges incident to  $x_1$  have distinct colors, 1 and 2, and that the edges  $x_1 y_2$  and  $x_2 y_1$  have the same color, 1. This implies that the edges  $y_1 z_i$  are oriented toward  $z_i$  and colored 3. This implies that the edges  $y_2 z_i$  are oriented toward  $z_i$  and colored 1. This finally implies that all the remaining edges incident to the  $z_i$ 's have the same color, which is not allowed. In the last drawing, we see a  $3-P_4$ -fa coloring of  $\mathcal{A}$  in which only one edge is oriented toward  $x_2$ .

This implies that there is not much flexibility for coloring the vertex gadget in Figure 4. Actually, in any  $3-P_4$ -fa coloring of the vertex gadget, the two edges incident to  $u$  have to be oriented toward  $u$  and so  $u$  is an inner vertex in exactly one forest. Indeed, if  $u x_1$  is oriented toward  $x_1$  and colored 3 then one copy of  $\mathcal{A}$ , say  $\mathcal{A}_1$ , has both edges  $x_1 y_1$  and  $x_1 y_2$  oriented from  $x_1$  to the other end. According to the possible  $3-P_4$ -fa colorings of  $\mathcal{A}_1$ , this implies that either  $x_1$  is an inner vertex in  $F_1$  and  $F_2$  and that  $x_2$  is an end in  $F_1$ , either that in  $\mathcal{A}_1$  both  $x_2 y_1$  and  $x_2 y_2$  are oriented toward  $x_2$ . In the first case the possible  $3-P_4$ -fa colorings of  $\mathcal{A}_2$  (the second copy of  $\mathcal{A}$ ) are such that  $x_2$  is an end in  $F_2$ . This implies that  $u x_2$  is colored 3 and oriented toward  $u$ , which is impossible since  $u x_1$  is also colored 3. In the second case  $\mathcal{A}_2$  should have only one edge oriented toward  $x_1$  and both edges incident to  $x_2$  oriented from  $x_2$  to the other end. Such  $3-P_4$ -fa coloring of  $\mathcal{A}_2$  would imply that  $x_2$  is an inner vertex in two forests which is impossible.

In any  $3-P_4$ -fa coloring of  $G'$ , since any vertex  $u \in V(G)$  is an inner vertex in exactly one forest, say  $F_1$ , the edges incident to  $u$  that belong to an edge gadget must be colored 1 and must be oriented toward the subdivision vertex. Thus the edge gadget forces two vertices  $u$  and  $v \in V(G)$  to be inner vertices in distinct forests. Now we show that  $G$  is 3-colorable iff  $G'$  is  $3-P_4$ -fa colorable.

Assume that  $G$  has a 3-coloring  $c$ , then for any vertex  $u \in V(G)$  we color its vertex gadget in  $G'$  so that  $u$  is an inner vertex in  $F_{c(u)}$ . We then extend this  $3-P_4$ -fa coloring to  $G'$ , which is possible since for any  $uv \in E(G)$  we have  $c(u) \neq c(v)$ . Conversely, suppose  $G'$  has a  $3-P_4$ -fa coloring, we color the vertices of  $G$  accordingly to the forest for which they are an inner vertex in the  $3-P_4$ -fa coloring of  $G'$ . This produces a 3-coloring of  $G$ .

### 4.3 Proof of Theorem 6.1

The graph  $\mathcal{A}$  depicted in Figure 6.(a) is such that in any of its  $3-S_n$ -fa colorings if none of the edges  $y_1 z_1$ ,  $y_1 z_2$ ,  $y_2 z_1$  and  $y_2 z_2$  uses a given color, say 3, then one of these edges is not oriented from  $y_i$  to  $z_j$ . Furthermore, there is such  $3-S_n$ -fa coloring of  $\mathcal{A}$  for which only one of these edges is unoriented.

The graph  $\mathcal{B}$  depicted in Figure 6.(b) is obtained using  $2n - 1$  copies of  $\mathcal{A}$ . The restrictions in the possible  $3-S_n$ -fa colorings of  $\mathcal{A}$  imply that in any  $3-S_n$ -fa coloring of  $\mathcal{B}$  one of the edges  $x_1 y_1$ ,  $x_1 y_2$ ,  $x_2 y_1$  and  $x_2 y_2$  is not oriented from




Fig. 6. (a) The graph  $\mathcal{A}$ , (b) The graph  $\mathcal{B}$  and (c) the vertex gadget of  $u$ .

$x_i$  to  $y_j$ . Furthermore, there is a  $3-S_n$ -fa coloring of  $\mathcal{B}$  where  $x_1y_1$ ,  $x_1y_2$  and  $x_2y_1$  are oriented towards  $y_j$  and respectively colored 1, 2 and 3; and where the edge  $x_2y_2$  is unoriented and colored 1.

The graph vertex gadget depicted in Figure 6.(c) is obtained using  $6(n - 1)$  copies of  $\mathcal{B}$ . The restrictions in the possible  $3-S_n$ -fa colorings of  $\mathcal{B}$  imply that in any of its  $3-S_n$ -fa colorings, the vertex  $u$  is an inner vertex in exactly one forest (the edges  $ux_1$  and  $ux_2$  are both oriented toward  $u$ ).

Given a planar graph  $G$ , we construct  $G'$  by adding to every vertex  $u$  of  $G$  the vertex gadget and by replacing every edge  $uv$  of  $G$  by a cycle  $(u, x_{uv}, v, y_{uv})$  where  $x_{uv}$  and  $y_{uv}$  are new vertices. The graph  $G'$  is clearly 2-degenerated, bipartite and planar. Now we prove that  $G$  is 3-colorable iff  $G'$  is  $3-S_n$ -fa colorable.

If  $G$  has a 3-coloring  $c$ , for each vertex  $u \in V(G)$  we  $3-S_n$ -fa color its gadget so that  $u$  is an inner vertex in  $F_{c(u)}$ . Then we orient the remaining edges incident to  $u$  from  $u$  to the other end and we color them  $c(u)$ . It is clear that for any edge  $uv \in E(G)$ , since  $c(u) \neq c(v)$  the cycle  $(u, x_{uv}, v, y_{uv})$  of  $G'$  is properly  $3-S_n$ -fa colored. So the graph  $G'$  is  $3-S_n$ -fa colorable.

Conversely, the restrictions in the possible  $3-S_n$ -fa colorings of a vertex gadget imply that if  $G'$  is  $3-S_n$ -fa colored any vertex  $u \in V(G)$  is an inner vertex in exactly one forest in  $G'$ . We define a 3-coloring  $c$  of  $G$  so that in  $G'$  any vertex  $u \in V(G)$  is an inner vertex in  $F_{c(u)}$ . Since for any cycle  $(u, x_{uv}, v, y_{uv})$  of  $G'$  the edges incident to  $u$  (resp.  $v$ ) are colored  $c(u)$  (resp.  $c(v)$ ) then for any edge  $uv \in E(G)$  we have  $c(u) \neq c(v)$ . So  $c$  is a 3-coloring of  $G$ .

#### 4.4 Proof of Theorem 6.2

Note that there is no  $2-S_n$ -fa coloring of the path  $(a, b, c, d)$  where the edges  $ab$  and  $cd$  are oriented toward  $b$  and  $c$  and have distinct colors. This implies that in  $\mathcal{A}$ , there is a forest, say  $F_1$ , such that both vertices  $a_i$  and  $b_j$  are inner vertices in  $F_1$ . This implies that in  $\mathcal{B}$ , the vertices  $a_i$  and  $a_j$  are respectively inner vertices in  $F_2$  and  $F_1$ . This implies that in  $\mathcal{C}$ , one of the  $a_i$ 's, say  $a_1$  (resp.




Fig. 7. The vertex gadget.

$a_2$ ), is an inner vertex in  $F_1$  (resp.  $F_2$ ) and that  $a_3$  is an inner vertex in both  $F_1$  and  $F_2$ . This implies that at least one of the edges  $u'a_i$  is unoriented. The possible colorings of  $\mathcal{B}$  also imply that in every  $2$ - $S_n$ - $fa$  coloring of  $\mathcal{D}$  where the edges  $ua_1$  and  $ua_2$  have the same color, one of these edges is unoriented. All this implies that in the vertex gadget (depicted in the right of Figure 7), the edge  $uu'$  is colored  $x \in \{1, 2\}$  and oriented toward  $u$  and that  $u$  is incident to  $n - 2$  unoriented edges colored  $3 - x$ .

Given a triangle-free planar graph  $G$ , we construct  $G'$  by adding to every vertex  $u \in V(G)$  the vertex gadget depicted in Figure 7, right, and by subdividing every edge of  $G$ . A  $2$ - $S_n$ - $fa$  coloring of the vertex gadget forces an original vertex of  $G$  to be an inner vertex in at most one forest, say  $F_1$ , and to be incident to at least  $n - 2$  unoriented edges of  $F_1$ . We consider now  $2$ - $S_n$ - $fa$  colorings of the edge gadget of an edge  $uv$  of  $G$ . If  $u$  and  $v$  are inner vertices in distinct forests, then we can  $2$ - $S_n$ - $fa$  color the edges of the edge gadget and orient them toward the subdivision vertex. If  $u$  and  $v$  are inner vertices in the forest  $F_1$ , then both edges of the edge gadget have to be unoriented edges colored 1. Thus  $u$  and  $v$  are now incident to  $n - 1$  unoriented edges of  $F_1$ . This shows that  $G$  has a  $2$ -subcoloring if and only if  $G'$  has a  $2$ - $S_n$ - $fa$  coloring.

## References

- [1] O. Borodin, A.V. Kostochka, J. Nešetřil, A. Raspaud, and E. Sopena. On the maximum average degree and the oriented chromatic number of a graph. *Discrete Math.*, 206:77–89, 1999.
- [2] O.V. Borodin, A.V. Kostochka, and D.R. Woodall. Acyclic colorings of planar graphs with large girth. *J. London Math. Soc.*, 60:344–352, 1999.
- [3] A. Brandstädt, V. B. Le, and J. P. Spinrad. *Graph classes: a survey*. SIAM Monographs on Discrete Mathematics and Applications. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.
- [4] H. Broersma, F. V. Fomin, J. Nešetřil, and G. J. Woeginger. More about subcolorings. *Computing*, 69(3):187–203, 2002.
- [5] J. Fiala, K. Jansen, V.B. Le, and E. Seidel. Graph subcolorings: Complexity and algorithms. *SIAM J. Discrete Math.*, 16(4):635–650, 2003.

- [6] J. Fiala and V.B. Le. The subchromatic index of graphs. *KAM Series*, 655, 2004.
- [7] M.R. Garey, D.S. Johnson, and L.J. Stockmeyer. Some simplified NP-complete graph problems. *Theor. Comput. Sci.*, 1(3):237–267, 1976.
- [8] J. Gimbel and C. Hartman. Subcolorings and the subchromatic number of a graph. *Discrete Math.*, 272:139–154, 2003.
- [9] D. Gonçalves. Caterpillar arboricity of planar graphs. *Discrete Math.*, 307(16):2112–2121, 2007.
- [10] A. Gyarfas and D.B. West. Multitrack interval graphs. *Congr. Numer.*, 139:109–116, 1995.
- [11] S.L. Hakimi, J. Mitchem, and E. Schmeichel. Star arboricity of graphs. *Discrete Math.*, 149:93–98, 1996.
- [12] A.M. Hobbs. Computing edge-toughness and fractional arboricity. *Contemp. Math.*, 89:89–106, 1989.
- [13] I. Holyer. The NP-completeness of edge coloring. *SIAM J. Comput.*, 10:718–720, 1981.
- [14] Alexandr V. Kostochka and Douglas B. West. Every outerplanar graph is the union of two interval graphs. In *Proceedings of the Thirtieth Southeastern International Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, FL, 1999)*, volume 139, pages 5–8, 1999.
- [15] C. St. J. A. Nash-Williams. Decomposition of finite graphs into forests. *J. London Math. Soc.*, 39:12, 1964.
- [16] E.R. Scheinerman and D.B. West. The interval number of a planar graph - three intervals suffice. *J. Combin. Theory Ser. B*, 35:224–239, 1983.
- [17] T.C. Shermer. On rectangle visibility graphs iii. external visibility and complexity. In *Proceedings of CCCG '96*, pages 234–239, 1996.
- [18] V. G. Vizing. On an estimate of the chromatic class of a  $p$ -graph. *Diskret. Analiz No.*, 3:25–30, 1964.