

# Separators & expansion

Jean-Florent Raymond

17<sup>th</sup> JCALM, Montpellier, March 2015

Talk based on: *Strongly sublinear separators and polynomial expansion*, Dvořák and Norin, 2015.

Talk based on: *Strongly sublinear separators and polynomial expansion*, Dvořák and Norin, 2015.

Theorem (Plotkin, Rao, and Smith, 1994)

$\forall G$ , *polynomial expansion*  $\Rightarrow$  *strongly sublinear separators*.

Theorem (Dvořák and Norin, 2015)


$\forall G$ , *strongly sublinear separators*  $\Rightarrow$  *polynomial expansion*.

Reminder of the previous talk:


- $r$ -minor of  $G$ : obtained by contracting disjoint balls of radius  $\leq r$  of a subgraph of  $G$ ;
- $\nabla_r(G) = \max \left\{ \frac{|E(G')|}{|V(G')|}, G' \text{ is a } r\text{-minor of } G \right\}$
- $G$  has **bounded expansion** if there is a function  $f$  such that

$$\forall r \in \mathbb{N}, \nabla_r(G) \leq f(r)$$

- $G$  has **polynomial expansion** if  $f$  is polynomial.


- separator  $(A, B)$ : 

# Separators

- separator  $(A, B)$ : 
- order of  $(A, B)$ :  $|A \cap B|$

# Separators

- separator  $(A, B)$ :




- order of  $(A, B)$ :  $|A \cap B|$

- balanced separator  $(A, B)$ :

$$|A \setminus B|, |B \setminus A| \leq \frac{2}{3} |V(G)|$$

# Separators


- separator  $(A, B)$ : 
- order of  $(A, B)$ :  $|A \cap B|$
- balanced separator  $(A, B)$ :

$$|A \setminus B|, |B \setminus A| \leq \frac{2}{3}|V(G)|$$

- $G$  has  $c \bullet^\delta$ -separators: every  $H \subseteq G$  has a balanced separator of order  $\leq c|V(H)|^\delta$ ;
- strongly sublinear separators:  $0 \leq \delta < 1$ .



# Separators

- separator  $(A, B)$ : 
- order of  $(A, B)$ :  $|A \cap B|$
- balanced separator  $(A, B)$ :

$$|A \setminus B|, |B \setminus A| \leq \frac{2}{3} |V(G)|$$

- $G$  has  $c \bullet^\delta$ -separators: every  $H \subseteq G$  has a balanced separator of order  $\leq c|V(H)|^\delta$ ;
- strongly sublinear separators:  $0 \leq \delta < 1$ .

Example: planar graphs have  $c \bullet^{\frac{1}{2}}$ -separators (for some constant  $c$ ).

# Why do we care about (small) separators?

- they give structural information;
- they are connected to several parameters (e.g. treewidth);
- they have algorithmic applications (divide and conquer).

# Why do we care about (small) separators?

- they give structural information;
- they are connected to several parameters (e.g. treewidth);
- they have algorithmic applications (divide and conquer).

Lemma (Robertson and Seymour, 1986)

*Any graph  $G$  has a balanced separator of order at most  $\mathbf{tw}(G) + 1$ .*

# Why do we care about (small) separators?

- they give structural information;
- they are connected to several parameters (e.g. treewidth);
- they have algorithmic applications (divide and conquer).

Lemma (Robertson and Seymour, 1986)

*Any graph  $G$  has a balanced separator of order at most  $\mathbf{tw}(G) + 1$ .*

Theorem (Dvořák and Norin, 2014)

$\forall c \geq 1, \beta \in [0, 1)$ , for every graph  $G$ ,

$G$  has  $c \bullet^\beta$  separators  $\Rightarrow \forall H \subseteq G, \mathbf{tw}(H) \leq 105c|V(H)|^\beta$ .

# Sublinear separators and polynomial expansion, once again

Recall:  $G$  has  $c \bullet^\delta$ -separators  $\equiv$  every  $H \subseteq G$  has a balanced separator of order  $\leq c|V(H)|^\delta$ .

# Sublinear separators and polynomial expansion, once again

Recall:  $G$  has  $c \bullet^\delta$ -separators  $\equiv$  every  $H \subseteq G$  has a balanced separator of order  $\leq c|V(H)|^\delta$ .

Theorem (Plotkin, Rao, and Smith, 1994)

$\forall d \geq 0$ , for every graph  $G$ ,

$G$  has expansion  $O(r^d) \Rightarrow G$  has  $c \bullet^{1 - \frac{1}{4d+3}}$ -separators  
(for some  $c \geq 1$ ).

# Sublinear separators and polynomial expansion, once again

Recall:  $G$  has  $c \bullet^\delta$ -separators  $\equiv$  every  $H \subseteq G$  has a balanced separator of order  $\leq c|V(H)|^\delta$ .

Theorem (Plotkin, Rao, and Smith, 1994)

$\forall d \geq 0$ , for every graph  $G$ ,

$G$  has expansion  $O(r^d) \Rightarrow G$  has  $c \bullet^{1 - \frac{1}{4d+3}}$ -separators  
(for some  $c \geq 1$ ).

Theorem (Dvořák and Norin, 2015)

$\forall c \geq 1, \delta \in (0, 1]$ , for every graph  $G$ ,

$G$  has  $c \bullet^{1-\delta}$ -separators  $\Rightarrow G$  has expansion  $O(r^{5/\delta^2})$ .

# A few words about expanders

$\alpha$ -expander  $G$ : at least  $\alpha|A|$  vertices of  $G \setminus A$  are adjacent to  $A$ ,  
for every  $A \subseteq V(G)$ ,  $|A| \leq |V(G)|/2$ .



# A few words about expanders

$\alpha$ -expander  $G$ : at least  $\alpha|A|$  vertices of  $G \setminus A$  are adjacent to  $A$ ,  
for every  $A \subseteq V(G)$ ,  $|A| \leq |V(G)|/2$ .

Lemma

$G$  is an  $\alpha$ -expander  $\Rightarrow \text{tw}(G) \geq \frac{\alpha}{3(1+\alpha)}$ .


# A few words about expanders

$\alpha$ -expander  $G$ : at least  $\alpha|A|$  vertices of  $G \setminus A$  are adjacent to  $A$ ,  
for every  $A \subseteq V(G)$ ,  $|A| \leq |V(G)|/2$ .

Lemma

$G$  is an  $\alpha$ -expander  $\Rightarrow \mathbf{tw}(G) \geq \frac{\alpha}{3(1+\alpha)}$ .

balanced separator  $(A, B)$  of order  $\leq \mathbf{tw}(G) + 1$ :



wlog  $|B \setminus A| \leq |V(H)|/2$


# A few words about expanders

$\alpha$ -expander  $G$ : at least  $\alpha|A|$  vertices of  $G \setminus A$  are adjacent to  $A$ ,  
for every  $A \subseteq V(G)$ ,  $|A| \leq |V(G)|/2$ .

Lemma

$G$  is an  $\alpha$ -expander  $\Rightarrow \mathbf{tw}(G) \geq \frac{\alpha}{3(1+\alpha)}$ .

balanced separator  $(A, B)$  of order  $\leq \mathbf{tw}(G) + 1$ :



wlog  $|B \setminus A| \leq |V(H)|/2$ , then  $|A \cap B| \geq \alpha|B \setminus A|$ .


# A few words about expanders

$\alpha$ -expander  $G$ : at least  $\alpha|A|$  vertices of  $G \setminus A$  are adjacent to  $A$ ,  
for every  $A \subseteq V(G)$ ,  $|A| \leq |V(G)|/2$ .

## Lemma

$G$  is an  $\alpha$ -expander  $\Rightarrow \mathbf{tw}(G) \geq \frac{\alpha}{3(1+\alpha)}$ .

balanced separator  $(A, B)$  of order  $\leq \mathbf{tw}(G) + 1$ :



wlog  $|B \setminus A| \leq |V(H)|/2$ , then  $|A \cap B| \geq \alpha|B \setminus A|$ .

we then use the fact that  $(A, B)$  is balanced and  
 $|A \cap B| \leq \mathbf{tw}(G) + 1$  to conclude.

# A few words about expanders

$\alpha$ -expander  $G$ : at least  $\alpha|A|$  vertices of  $G \setminus A$  are adjacent to  $A$ ,  
for every  $A \subseteq V(G)$ ,  $|A| \leq |V(G)|/2$ .

Lemma

$G$  is an  $\alpha$ -expander  $\Rightarrow \text{tw}(G) \geq \frac{\alpha}{3(1+\alpha)}$ .

Lemma (Dvořák and Norin, 2015)

For every  $\varepsilon \in (0, 1]$  and every  $t$  large enough,

if  $G$  has  $\geq f(\varepsilon)t^4|V(G)|^{1+\varepsilon}$  edges

then  $\exists H \subseteq G$  subcubic s.t.  $|V(H)| \leq f'(\varepsilon)t$  and  $\text{tw}(G) \geq \frac{t}{25}$ .

# Outline of the proof

Let  $c \geq 1$  and  $\delta \in (0, 1]$ .

Goal:  $G$  has  $c \bullet^{1-\delta}$ -separators  $\Rightarrow G$  has expansion  $O(r^{5/\delta^2})$ .

# Outline of the proof

Let  $c \geq 1$  and  $\delta \in (0, 1]$ .

Goal:  $G$  has  $c \bullet^{1-\delta}$ -separators  $\Rightarrow G$  has expansion  $O(r^{5/\delta^2})$ .

Let us assume that:

1.  $\forall r, \forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-5\delta/6}$ -separators;

# Outline of the proof

Let  $c \geq 1$  and  $\delta \in (0, 1]$ .

Goal:  $G$  has  $c \bullet^{1-\delta}$ -separators  $\Rightarrow G$  has expansion  $O(r^{5/\delta^2})$ .

Let us assume that:

1.  $\forall r, \forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-5\delta/6}$ -separators;
2.  $\forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then  $|E(G)| \leq O_\delta((c \log^3 c)^{1/\delta}) \cdot |V(H)|$ .



# Outline of the proof

Let  $c \geq 1$  and  $\delta \in (0, 1]$ .

Goal:  $G$  has  $c \bullet^{1-\delta}$ -separators  $\Rightarrow G$  has expansion  $O(r^{5/\delta^2})$ .

Let us assume that:

1.  $\forall r, \forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-5\delta/6}$ -separators;
2.  $\forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then  $|E(G)| \leq O_\delta((c \log^3 c)^{1/\delta}) \cdot |V(H)|$ .

We apply (2.) to an  $r$ -minor  $H$  of  $G$ :  $|E(H)| \leq O\left(r^{5/\delta^2}\right) \cdot |V(H)|$ .

# Outline of the proof

Let  $c \geq 1$  and  $\delta \in (0, 1]$ .

Goal:  $G$  has  $c \bullet^{1-\delta}$ -separators  $\Rightarrow G$  has expansion  $O(r^{5/\delta^2})$ .

Let us assume that:

1.  $\forall r, \forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-5\delta/6}$ -separators;
2.  $\forall G$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then  $|E(G)| \leq O_\delta((c \log^3 c)^{1/\delta}) \cdot |V(H)|$ .

We apply (2.) to an  $r$ -minor  $H$  of  $G$ :  $|E(H)| \leq O(r^{5/\delta^2}) \cdot |V(H)|$ .

Hence  $\nabla_r(G) = O(r^{5/\delta^2})$ .

$G$  has polynomial expansion.

# Strongly sublinear separators force low density

## Lemma (2.)

For every  $G$  graph, if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then  $|E(G)| \leq O_\delta ((c \log^3 c)^{1/\delta}) \cdot |V(G)|$

# Strongly sublinear separators force low density

## Lemma (2.)

For every  $G$  graph, if  $G$  has  $c^{1-\delta}$ -separators,  
then  $|E(G)| \leq O_\delta((c \log^3 c)^{1/\delta}) \cdot |V(G)|$

Goal: show that  $|E(G)| \leq f_\delta(c) |V(G)| \left(1 - \frac{1}{\log |V(G)|}\right)$ .  
for some  $f_\delta(c) = O_\delta((c \log^3 c)^{1/\delta})$

# Strongly sublinear separators force low density


## Lemma (2.)

For every  $G$  graph, if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then  $|E(G)| \leq O_\delta((c \log^3 c)^{1/\delta}) \cdot |V(G)|$

Goal: show that  $|E(G)| \leq f_\delta(c) |V(G)| \left(1 - \frac{1}{\log |V(G)|}\right)$ .

for some  $f_\delta(c) = O_\delta((c \log^3 c)^{1/\delta})$

Let  $(A, B)$  be a balanced separator of order  $\leq c |V(G)|^{1-\delta}$ .



# Strongly sublinear separators force low density


## Lemma (2.)

For every  $G$  graph, if  $G$  has  $c \bullet^{1-\delta}$ -separators, then  $|E(G)| \leq O_\delta((c \log^3 c)^{1/\delta}) \cdot |V(G)|$

Goal: show that  $|E(G)| \leq f_\delta(c) |V(G)| \left(1 - \frac{1}{\log |V(G)|}\right)$ .

for some  $f_\delta(c) = O_\delta((c \log^3 c)^{1/\delta})$

Let  $(A, B)$  be a balanced separator of order  $\leq c |V(G)|^{1-\delta}$ .



$$\begin{aligned} |E(G)| &\leq |E(G[A])| + |E(G[B])| \\ &\leq f_\delta(c) |A| \left(1 - \frac{1}{\log |A|}\right) + f_\delta(c) |B| \left(1 - \frac{1}{\log |B|}\right) \\ &\leq f_\delta(c) n \left(1 - \frac{1}{\log n}\right) \end{aligned}$$

# The densities of bounded-depth minors

Before proving

Lemma (1.)

For every  $r \geq 1$ ,

if  $G$  has  $c \bullet^{1-\delta}$ -separators,

then every  $r$ -minor of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

# The densities of bounded-depth minors

Before proving

## Lemma (1.)

For every  $r \geq 1$ ,  
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

we show

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta}) |V(H)|^{1+\varepsilon}$  edges.



# The densities of bounded-depth minors

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

# The densities of bounded-depth minors

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

1. suppose that  $|E(H)| = \Omega_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  (by contradiction).

# The densities of bounded-depth minors

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

1. suppose that  $|E(H)| = \Omega_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  (by contradiction).
2.  $\exists F \subseteq H$  subcubic s.t  $|V(F)|$  is *small* and  $\mathbf{tw}(F)$  is *large*  
(using expanders).

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

1. suppose that  $|E(H)| = \Omega_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  (by contradiction).
2.  $\exists F \subseteq H$  subcubic s.t.  $|V(F)|$  is *small* and  $\mathbf{tw}(F)$  is *large*  
(using expanders).
3.  $\exists F' \subseteq G$  subcubic s.t.  $F \leq_m F'$  and we have  $\mathbf{tw}(F) \leq \mathbf{tw}(F')$ .

# The densities of bounded-depth minors

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

1. suppose that  $|E(H)| = \Omega_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  (by contradiction).
2.  $\exists F \subseteq H$  subcubic s.t.  $|V(F)|$  is *small* and  $\mathbf{tw}(F)$  is *large*  
(using expanders).
3.  $\exists F' \subseteq G$  subcubic s.t.  $F \leq_m F'$  and we have  $\mathbf{tw}(F) \leq \mathbf{tw}(F')$ .
4.  $F'$  has  $c \bullet^{1-\delta}$ -separators.

# The densities of bounded-depth minors

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

1. suppose that  $|E(H)| = \Omega_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  (by contradiction).
2.  $\exists F \subseteq H$  subcubic s.t.  $|V(F)|$  is *small* and  $\mathbf{tw}(F)$  is *large*  
(using expanders).
3.  $\exists F' \subseteq G$  subcubic s.t.  $F \leq_m F'$  and we have  $\mathbf{tw}(F) \leq \mathbf{tw}(F')$ .
4.  $F'$  has  $c \bullet^{1-\delta}$ -separators.
5.  $\mathbf{tw}(F') \leq 105c|V(F')|^{1-\delta}$  (*small*).

# The densities of bounded-depth minors

## Lemma

For every  $r \geq 1, \varepsilon \in (0, 1]$   
if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  edges.

Goal:  $|E(H)| = O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

1. suppose that  $|E(H)| = \Omega_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$  (by contradiction).
2.  $\exists F \subseteq H$  subcubic s.t.  $|V(F)|$  is *small* and  $\mathbf{tw}(F)$  is *large*  
(using expanders).
3.  $\exists F' \subseteq G$  subcubic s.t.  $F \leq_m F'$  and we have  $\mathbf{tw}(F) \leq \mathbf{tw}(F')$ .
4.  $F'$  has  $c \bullet^{1-\delta}$ -separators.
5.  $\mathbf{tw}(F') \leq 105c|V(F')|^{1-\delta}$  (*small*).
6.  $\mathbf{tw}(F) \leq 105c|V(F')|^{1-\delta}$  and  $\mathbf{tw}(F)$  is *large*: contradiction.



## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators,  
then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.


1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2. Let us find a small subgraph of  $G$  containing  $H$  as a minor:




## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2. Let us find a small subgraph of  $G$  containing  $H$  as a minor:




## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2. Let us find a small subgraph of  $G$  containing  $H$  as a minor:



Every edge of  $F$


- meets two balls;
- contributes to  $2r$  vertices.

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2. Let us find a small subgraph of  $G$  containing  $H$  as a minor:



Every edge of  $F$

- meets two balls;
- contributes to  $2r$  vertices.

$H$  is a minor of some  $H' \subseteq G$  s.t.  
 $|V(H')| \leq 2r|E(H)| + |V(H)|$ .

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2.  $H$  is a minor of  $H' \subseteq G$  with  $|V(H')| \leq 2r|E(H)| + |V(H)|$ .



## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2.  $H$  is a minor of  $H' \subseteq G$  with  $|V(H')| \leq 2r|E(H)| + |V(H)|$ .
3.  $H'$  has  $c \bullet^{1-\delta}$ -separators.

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2.  $H$  is a minor of  $H' \subseteq G$  with  $|V(H')| \leq 2r|E(H)| + |V(H)|$ .
3.  $H'$  has  $c \bullet^{1-\delta}$ -separators.
4.  $\mathbf{tw}(H) \leq \mathbf{tw}(H') \leq 105c|V(H')|^{1-\delta}$ .

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2.  $H$  is a minor of  $H' \subseteq G$  with  $|V(H')| \leq 2r|E(H)| + |V(H)|$ .
3.  $H'$  has  $c \bullet^{1-\delta}$ -separators.
4.  $\mathbf{tw}(H) \leq \mathbf{tw}(H') \leq 105c|V(H')|^{1-\delta}$ .
5.  $H$  has a balanced separator of order  $\leq \mathbf{tw}(H) + 1$ .

## Lemma (1.)

For every  $r \geq 1$ , if  $G$  has  $c \bullet^{1-\delta}$ -separators, then every  $r$ -minor  $H$  of  $G$  has  $O_{c,\delta}(r^{4/\delta}) \bullet^{1-\frac{5\delta}{6}}$ -separators.

Goal: show that  $H$  has a sublinear balanced separator.

1. we proved that  $\forall \varepsilon \in (0, 1]$ ,  $|E(H)| \leq O_{c,\delta,\varepsilon}(r^{4/\delta})|V(H)|^{1+\varepsilon}$ .
2.  $H$  is a minor of  $H' \subseteq G$  with  $|V(H')| \leq 2r|E(H)| + |V(H)|$ .
3.  $H'$  has  $c \bullet^{1-\delta}$ -separators.
4.  $\mathbf{tw}(H) \leq \mathbf{tw}(H') \leq 105c|V(H')|^{1-\delta}$ .
5.  $H$  has a balanced separator of order  $\leq \mathbf{tw}(H) + 1$ .
6.  $H$  has a balanced separator of order  $O_{c,\delta}(r^{4/\delta})|V(H)|^{1-\frac{5\delta}{6}}$ .

We proved that

strongly sublinear separators  $\Rightarrow$  polynomial expansion.

We proved that

**strongly sublinear separators**  $\Rightarrow$  **polynomial expansion**.

In some sense, **strongly sublinear separators** and **polynomial expansion** are equivalent notions.

We proved that

**strongly sublinear separators**  $\Rightarrow$  **polynomial expansion**.

In some sense, **strongly sublinear separators** and **polynomial expansion** are equivalent notions.

A class  $\mathcal{C}$  has **strongly sublinear separators** if  $\exists c \geq 1, \delta \in (0, 1]$  such that every  $G \in \mathcal{C}$  has balanced separators of order  $\leq c|V(G)|^{1-\delta}$ .

We proved that

**strongly sublinear separators**  $\Rightarrow$  **polynomial expansion**.

In some sense, **strongly sublinear separators** and **polynomial expansion** are equivalent notions.

A class  $\mathcal{C}$  has **strongly sublinear separators** if  $\exists c \geq 1, \delta \in (0, 1]$  such that every  $G \in \mathcal{C}$  has balanced separators of order  $\leq c|V(G)|^{1-\delta}$ .

## Consequence of the main result

Let  $\mathcal{C}$  be a subgraph-closed class.

**$\mathcal{C}$  has strongly sublinear separators** iff  **$\mathcal{C}$  has polynomial expansion**.



We proved that

**strongly sublinear separators**  $\Rightarrow$  **polynomial expansion**.

In some sense, **strongly sublinear separators** and **polynomial expansion** are equivalent notions.

A class  $\mathcal{C}$  has **strongly sublinear separators** if  $\exists c \geq 1, \delta \in (0, 1]$  such that every  $G \in \mathcal{C}$  has balanced separators of order  $\leq c|V(G)|^{1-\delta}$ .

## Consequence of the main result

Let  $\mathcal{C}$  be a subgraph-closed class.

**$\mathcal{C}$  has strongly sublinear separators** iff  **$\mathcal{C}$  has polynomial expansion**.

Rk: if  **$\mathcal{C}$  is subgraph-closed** then every  $G \in \mathcal{C}$  has  **$c \bullet^{1-\delta}$**  separators.

## Conjecture

$\exists k > 0, \forall c \geq 1, \delta \in (0, 1],$

if a graph  $G$  has  $c \bullet^{1-\delta}$  separators, then its expansion is  $O(r^{k/\delta})$ .

Currently:  $O(r^{5/\delta^2})$ .

## Conjecture

$\exists k > 0, \forall c \geq 1, \delta \in (0, 1]$ ,  
if a graph  $G$  has  $c \bullet^{1-\delta}$  separators, then its expansion is  $O(r^{k/\delta})$ .

Currently:  $O(r^{5/\delta^2})$ .

**Thank you for your attention!**