

Turán numbers and anti-Ramsey numbers for short cycles in complete 3-partite graphs

Ervin Győri — Alfréd Rényi Institute of Mathematics, Budapest

Chunqiu Fang — School of Mathematical Sciences, University of Science and Technology of China, Hefei

Chuanqi Xiao — Central European University, Budapest

Jimeng Xiao — School of Mathematics and Statistics, Northwestern Polytechnical University, Xi'an

Abstract

We call a 4-cycle in K_{n_1, n_2, n_3} multipartite, denoted by C_4^{multi} , if it contains at least one vertex in each part of K_{n_1, n_2, n_3} . The Turán number $ex(K_{n_1, n_2, n_3}, C_4^{\text{multi}})$ (respectively, $ex(K_{n_1, n_2, n_3}, \{C_3, C_4^{\text{multi}}\})$) is the maximum number of edges in a graph $G \subseteq K_{n_1, n_2, n_3}$ such that G contains no C_4^{multi} (respectively, G contains neither C_3 nor C_4^{multi}). We call an edge-colored C_4^{multi} rainbow if all four edges of it have different colors. The anti-Ramsey number $ar(K_{n_1, n_2, n_3}, C_4^{\text{multi}})$ is the maximum number of colors in an edge-colored K_{n_1, n_2, n_3} with no rainbow C_4^{multi} . In this paper, we determine that $ex(K_{n_1, n_2, n_3}, C_4^{\text{multi}}) = n_1 n_2 + 2n_3$ and $ar(K_{n_1, n_2, n_3}, C_4^{\text{multi}}) = ex(K_{n_1, n_2, n_3}, \{C_3, C_4^{\text{multi}}\}) + 1 = n_1 n_2 + n_3 + 1$, where $n_1 \geq n_2 \geq n_3 \geq 1$.

1 Introduction

After extended study of Ramsey problems - finding monochromatic subgraphs - anti-Ramsey questions were raised and discussed first by Erdős, Simonovits and Sós [7]. A subgraph of an edge-colored graph is *rainbow*, if all of its edges have different colors. For graphs G and H , the anti-Ramsey number $ar(G, H)$ is the maximum number of colors in an edge-colored G with no rainbow copy of H . Erdős, Simonovits and Sós [7] first studied the anti-Ramsey number in the case when the host graph G is a complete graph K_n and showed the close relationship between it and the Turán number. Since then, the anti-Ramsey numbers for various special graph classes - e.g., paths, cycles, cliques and matchings - in the complete graph have been determined, see e.g. [1, 5, 9, 12, 13, 15]. Later, many different graphs have been considered as the host graph G : complete bipartite or r -partite graphs, complete hypergraphs, complete split graphs and triangulations, etc. In this paper, we consider the anti-Ramsey number of C_4^{multi} in the complete 3-partite graphs.

We consider only nonempty simple graphs. Let G be such a graph, the vertex and edge set of G is denoted by $V(G)$ and $E(G)$, the number of vertices and edges in G by $\nu(G)$ and $e(G)$, respectively. We denote the neighborhood of v in G by $N_G(v)$, and the degree of a vertex v in G by $d_G(v)$, the size of $N_G(v)$. Let U_1, U_2 be vertex sets, denote by $e_G(U_1, U_2)$ the number of edges between U_1 and U_2 in G . We write $d(v)$ instead of $d_G(v)$, $N(v)$ instead of $N_G(v)$ and $e(U_1, U_2)$ instead of $e_G(U_1, U_2)$ if the underlying graph G is clear.

Given a graph family \mathcal{F} , we call a graph H an \mathcal{F} -free graph, if H contains no graph in \mathcal{F} as a subgraph. The Turán number $ex(G, \mathcal{F})$ for a graph family \mathcal{F} in G is the maximum number of edges in a graph $H \subseteq G$ which is \mathcal{F} -free. If $\mathcal{F} = \{F\}$, then we denote $ex(G, \mathcal{F})$ by $ex(G, F)$.

An old result of Bollobás, Erdős and Szemerédi [3] showed that $ex(K_{n_1, n_2, n_3}, C_3) = n_1 n_2 + n_1 n_3$ for $n_1 \geq n_2 \geq n_3 \geq 1$ (also see [4, 2, 6]). Lv, Lu and Fang [10, 11] constructed balanced 3-partite graphs which are C_4 -free and $\{C_3, C_4\}$ -free respectively and showed that $ex(K_{n, n, n}, C_4) = (\frac{3}{\sqrt{2}} + o(1))n^{3/2}$ and $ex(K_{n, n, n}, \{C_3, C_4\}) \geq (\sqrt{3} + o(1))n^{3/2}$.

For further discussion, we need the definitions of the multipartite subgraphs and a function $f(n_1, n_2, \dots, n_r)$.

Definition 1. [8] Let $r \geq 3$ and G be an r -partite graph with vertex partition V_1, V_2, \dots, V_r , we call a subgraph H of G multipartite, if there are at least three distinct parts V_i, V_j, V_k such that $V(H) \cap V_i \neq \emptyset, V(H) \cap V_j \neq \emptyset$ and $V(H) \cap V_k \neq \emptyset$. In particular, we denote a multipartite H by H^{multi} .

Fang, Győri, Li and Xiao [8] recently showed that if $G \subseteq K_{n_1, n_2, \dots, n_r}$ and $e(G) \geq f(n_1, n_2, \dots, n_r) + 1$, then G contains a multipartite cycle. Furthermore, they proposed the following conjecture.

Conjecture 1. [8] For $r \geq 3$ and $n_1 \geq n_2 \geq \dots \geq n_r \geq 1$, if $G \subset K_{n_1, n_2, \dots, n_r}$ and $e(G) \geq f(n_1, n_2, \dots, n_r) + 1$, then G contains a multipartite cycle C^{multi} of length at most $\frac{3}{2}r$.

In this paper, we study the Turán numbers of C_4^{multi} and $\{C_3, C_4^{multi}\}$ in the complete 3-partite graphs and obtain the following results.

Theorem 1. For $n_1 \geq n_2 \geq n_3 \geq 1$, $ex(K_{n_1, n_2, n_3}, C_4^{multi}) = n_1 n_2 + 2n_3$.

Lower bound: Let $G \subseteq K_{n_1, n_2, n_3}$ be a graph, such that V_1 and V_2 are completely joined, V_1 (respectively, V_2) and V_3 are joined by an n_3 -matching. Clearly, G is C_4^{multi} -free and $e(G) = n_1 n_2 + 2n_3$. Therefore, $ex(K_{n_1, n_2, n_3}, C_4^{multi}) \geq n_1 n_2 + 2n_3$.

Theorem 2. For $n_1 \geq n_2 \geq n_3 \geq 1$, $ex(K_{n_1, n_2, n_3}, \{C_3, C_4^{multi}\}) = n_1 n_2 + n_3$.

Lower bound: Let $G \subseteq K_{n_1, n_2, n_3}$ be a graph, such that V_1 and V_2 are completely joined, V_1 and V_3 are joined by an n_3 -matching and there is no edge between V_2 and V_3 . Clearly, G is $\{C_3, C_4^{multi}\}$ -free and $e(G) = n_1 n_2 + n_3$. Therefore, $ex(K_{n_1, n_2, n_3}, \{C_3, C_4^{multi}\}) \geq n_1 n_2 + n_3$.

Notice that Theorem 2 confirms Conjecture 1 for the case when $r = 3$.

Theorem 3. For $n_1 \geq n_2 \geq n_3 \geq 1$, $ar(K_{n_1, n_2, n_3}, C_4^{multi}) = n_1 n_2 + n_3 + 1$.

Lower bound: We color the edges of K_{n_1, n_2, n_3} as follows. First, color all edges between V_1 and V_2 rainbow. Second, for each vertex $v \in V_3$, color all the edges between v and V_1 with one new distinct color. Finally, we assign a new color to all edges between V_2 and V_3 . In such way, we use exactly $n_1 n_2 + n_3 + 1$ colors, and there is no rainbow C_4^{multi} .

The following lemma plays an important role in our proof.

Lemma 1. Let G be a 3-partite graph with vertex partition X, Y and Z , such that for all $x \in X$, $N(x) \cap Y \neq \emptyset$ and $N(x) \cap Z \neq \emptyset$.

- (i) If G is C_4^{multi} -free, then $e(G) \leq |Y||Z| + 2|X|$;
- (ii) If G is $\{C_3, C_4^{multi}\}$ -free, then $e(G) \leq |Y||Z| + |X|$.

Proof. (i) Since G is C_4^{multi} -free, $G[N(x)]$ is $K_{1,2}$ -free for each $x \in X$. Therefore,

$$e(G[N(x)]) = e\left(N(x) \cap Y, N(x) \cap Z\right) \leq \min\left\{|N(x) \cap Y|, |N(x) \cap Z|\right\}. \quad (1)$$

For $x \in X$, we let e_x be the number of missing edges of G between $N(x) \cap Y$ and $N(x) \cap Z$. By (1), we have

$$\begin{aligned} e_x &= |N(x) \cap Y| \cdot |N(x) \cap Z| - e\left(N(x) \cap Y, N(x) \cap Z\right) \\ &\geq |N(x) \cap Y| \cdot |N(x) \cap Z| - \min\left\{|N(x) \cap Y|, |N(x) \cap Z|\right\} \\ &\geq |N(x) \cap Y| + |N(x) \cap Z| - 2, \end{aligned} \quad (2)$$

where the last inequality holds since $|N(x) \cap Y| \geq 1$ and $|N(x) \cap Z| \geq 1$ for all $x \in X$.

By (2), we get

$$\sum_{x \in X} e_x \geq \sum_{x \in X} \left(|N(x) \cap Y| + |N(x) \cap Z| - 2\right) = e(X, Y) + e(X, Z) - 2|X|. \quad (3)$$

Notice that for any two distinct vertices $x_1, x_2 \in X$, they cannot have common neighbors in both Y and Z at the same time, otherwise we find a copy of C_4^{multi} in G . Thus each missing edge between Y and Z is calculated at most once in the sum $\sum_{x \in X} e_x$. Hence the number of missing edges between Y and Z is at least $\sum_{x \in X} e_x$. Then we have

$$e(Y, Z) \leq |Y||Z| - \sum_{x \in X} e_x \leq |Y||Z| - (e(X, Y) + e(X, Z) - 2|X|). \quad (4)$$

By (4), we get

$$e(G) = e(X, Y) + e(X, Z) + e(Y, Z) \leq |Y||Z| + 2|X|.$$

(ii) Since G is C_3 -free, for each $x \in X$,

$$e\left(N(x) \cap Y, N(x) \cap Z\right) = 0. \quad (5)$$

Since for each $x \in X$, $|N(x) \cap Y| \geq 1$ and $|N(x) \cap Z| \geq 1$ hold, by (5), the number of missing edges between $N(x) \cap Y$ and $N(x) \cap Z$ is $|N(x) \cap Y| \cdot |N(x) \cap Z|$. Notice that for any two distinct vertices $x_1, x_2 \in X$, they cannot have common neighbors in both Y and Z at the same time, otherwise we find a copy of C_4^{multi} in G . Hence, the number of missing edges between Y and Z is at least $\sum_{x \in X} |N(x) \cap Y| \cdot |N(x) \cap Z|$. Thus,

$$\begin{aligned} e(Y, Z) &\leq |Y||Z| - \sum_{x \in X} |N(x) \cap Y| \cdot |N(x) \cap Z| \\ &\leq |Y||Z| - \sum_{x \in X} (|N(x) \cap Y| + |N(x) \cap Z| - 1) \\ &= |Y||Z| + |X| - e(X, Y) - e(X, Z), \end{aligned} \quad (6)$$

the second inequality holds since $|N(x) \cap Y| \geq 1$ and $|N(x) \cap Z| \geq 1$ for $x \in X$.

By (6), we have $e(G) = e(Y, Z) + e(X, Y) + e(X, Z) \leq |Y||Z| + |X|$. \square

The proof of the general case when the conditions of this lemma do not hold is several pages, we cannot include it in this extended abstract.

Acknowledgment

The research of the first author was supported in part by Anhui Initiative in Quantum Information Technologies grant AHY150200 and NSFC (No. 11771247). The research of second and third authors was partially supported by the National Research, Development and Innovation Office NKFIH, grants K132696, K116769, SNN-135643. The research of the fourth author was supported in part by NSFC (No. 12071370).

References

- [1] N. Alon, On a Conjecture of Erdős, Simonovits and Sós Concerning Anti-Ramsey Theorems. *J. Graph Theory* 7 (1) (1983) 91-94.
- [2] P. Bennett, S. English and M. Talanda-Fisher, Weighted Turán problems with applications, *Discrete Math.* 342 (8) (2019) 2165-2172.
- [3] B. Bollobás, *Extremal Graph Theory*, Academic Press, New York, 1978.
- [4] B. Bollobás, P. Erdős, E. Szemerédi, On complete subgraphs of r -chromatic graphs, *Discrete Math.* 13 (2) (1975) 97-107.
- [5] H. Chen, X. Li, J. Tu, Complete solution for the rainbow numbers of matchings, *Discrete math.* 309 (10) (2009) 3370-3380.
- [6] J. De Silva, K. Heysse, A. Kapilow, A. Schenfisch, M. Young, Turán numbers of vertex-disjoint cliques in r -partite graphs, *Discrete Math.* 341 (2) (2018), 492-496.
- [7] P. Erdős, M. Simonovits, V. T. Sós, Anti-Ramsey theorems, Infinite and finite sets, in: *Colloq. Math. Soc. Janos Bolyai*. North Holland, Vol. 10 (1975) 633-643.
- [8] C. Fang, E. Győri, B. Li, J. Xiao, The anti-Ramsey number of C_3 and C_4 in the complete r -partite graphs, arXiv:2007.06003.
- [9] R. Haas, M. Young, The anti-Ramsey number of perfect matching, *Discrete Math.* 312 (5) (2012) 993-937.
- [10] Z. Lv, M. Lu, C. Fang, A note on 3-partite graphs without 4-cycles, *J. Combin. Des.* 28 (10) (2020) 753-757.
- [11] Z. Lv, M. Lu, C. Fang, A note on 3-partite graphs without 3-cycles or 4-cycles, submitted.
- [12] J.J. Montellano-Ballesteros, V. Neumann-Lara, An anti-Ramsey theorem on cycles, *Graphs Combin.* 21 (3) (2005) 343-354.
- [13] J. J. Montellano-Ballesteros, V. Neumann-Lara, An anti-Ramsey theorem, *Combinatorica* 22 (3) (2002) 445-449.
- [14] L. Özkahya, M. Young, Anti-Ramsey number of matchings in hypergraphs, *Discrete Math.* 313 (20) (2013) 2359-2364.
- [15] I. Schiermeyer, Rainbow numbers for matchings and complete graphs, *Discrete Math* 286 (1-2) (2004) 157-162.