

# A bound on the dissociation number

Felix Bock — Ulm University, Germany

Johannes Pardey — Ulm University, Germany

Lucia D. Penso — Ulm University, Germany

Dieter Rautenbach — Ulm University, Germany

## Abstract

The dissociation number  $\text{diss}(G)$  of a graph  $G$  is the maximum order of a set of vertices of  $G$  inducing a subgraph that is of maximum degree at most 1. Computing the dissociation number of a given graph is algorithmically hard even when restricted to subcubic bipartite graphs. For a graph  $G$  with  $n$  vertices,  $m$  edges,  $k$  components, and  $c_1$  induced cycles of length 1 modulo 3, we show  $\text{diss}(G) \geq n - \frac{1}{3}(m + k + c_1)$ . Furthermore, we characterize the extremal graphs in which every two cycles are vertex-disjoint.

**Keywords:** Dissociation set, dissociation number, cactus

## 1 Introduction

We consider finite, simple, and undirected graphs, and use standard terminology. A set  $D$  of vertices of a graph  $G$  is a *dissociation set* in  $G$  if the subgraph  $G[D]$  of  $G$  induced by  $D$  has maximum degree at most 1. The *dissociation number*  $\text{diss}(G)$  of  $G$  is the maximum order of a dissociation set in  $G$ . The dissociation number is algorithmically hard even when restricted, for instance, to subcubic bipartite graphs [3, 9, 12]. Fast exact algorithms [8], (randomized) approximation algorithms [7, 8], and fixed parameter tractability [11] have been studied for this parameter or its dual, the *3-path (vertex) cover number*. Several lower bounds on the dissociation number were proposed: If  $G$  is a graph of order  $n$  and size  $m$ , then

$$\text{diss}(G) \geq \begin{cases} \left\lceil \frac{n}{\frac{\Delta+1}{2}} \right\rceil & , \text{ if } G \text{ has maximum degree } \Delta \text{ [4],} \\ \frac{4}{3} \sum_{u \in V(G)} \frac{1}{d_G(u)+1} & , \text{ if } G \text{ has no isolated vertex [4],} \\ \sum_{u \in V(G)} \frac{1}{d_G(u)+1} + \sum_{uv \in E(G)} \binom{|N_G[u] \cup N_G[v]|}{2}^{-1} & , \text{ [6],} \\ \frac{n}{2} & , \text{ if } G \text{ is outerplanar [4]} \\ \frac{2n}{3} & , \text{ if } G \text{ is a tree [4],} \\ \frac{2n}{k+2} - \frac{m}{(k+1)(k+2)} & , \text{ if } k = \left\lceil \frac{m}{n} \right\rceil - 1 \text{ [5], and} \\ \frac{2n}{3} - \frac{m}{6} & , \text{ [4].} \end{cases} \quad (1)$$

The results in the present papers were inspired by bounds in (1).

Our main result is the following.

**Theorem 1.** *If  $G$  is a graph with  $n$  vertices,  $m$  edges,  $k$  components, and  $c_1$  induced cycles of length 1 modulo 3, then*

$$\text{diss}(G) \geq n - \frac{1}{3}(m + k + c_1). \quad (2)$$

Theorem 1 generalizes the lower bound  $2n/3$  for trees of order  $n$  in (1), strengthens the general lower bound  $\frac{2n}{3} - \frac{m}{6}$  in (1) for many graphs, and almost implies the lower bound  $n/2$  for subcubic graphs of order  $n$ , which follows from the first bound in (1). In the proof of Theorem 1, graphs in which all cycles are pairwise vertex-disjoint play an essential role. We call such graphs *cycle-disjoint*; their components are restricted cactus graphs, where a *cactus* is a connected graph in which every block is either a  $K_2$  or a cycle. As a step towards the understanding of all extremal graphs for Theorem 1, we consider the extremal cycle-disjoint graphs in more detail. We propose three extension operations  $(O_1)$ ,  $(O_2)$ , and  $(O_3)$  applicable to a given graph  $G'$ , attaching a  $P_3$  or a cycle of length not 0 modulo 3 by a bridge to  $G'$ , illustrated in Figure 1. It is easy to see that applying one of these operations to a graph that satisfies (2) with equality yields a graph that satisfies (2) with equality. Since  $P_3$  and the cycles of lengths not 0 modulo 3 satisfy (2) with equality, this already allows to construct quite a rich family of extremal graphs, yet not all of them.




Figure 1: Operations constructing an extremal graph from a smaller extremal graph  $G'$ .

The two operations  $(O_1)$  and  $(O_2)$  are sufficient for the constructive characterization of all trees  $T$  of order  $n$  with  $\text{diss}(T) = 2n/3$ , that is, of all trees that are extremal for the bound from [4] stated in (1). Let  $\mathcal{T}$  be the set of all trees that arise from  $P_3$  by repeated applications of the two operations  $(O_1)$  and  $(O_2)$ , attaching a new  $P_3$  by a bridge to trees in  $\mathcal{T}$ .

**Theorem 2.** *For a tree  $T$  of order  $n$ , the following statements are equivalent.*

- (a)  $\text{diss}(T) = \frac{2n}{3}$ .
- (b)  $T \in \mathcal{T}$ .
- (c)  $n \equiv 0 \pmod{3}$ , and, for every vertex  $y$  of  $T$ , at most two components of  $T - y$  have order not 0 modulo 3.

Next to the three simple operations illustrated in Figure 1, we introduce one slightly more complicated operation involving so-called (*very*) *good* *spiked cycles*: For positive integers  $\ell$  and  $k$  with  $\ell \geq \max\{3, k\}$ , and indices  $i_1, \dots, i_k \in [\ell]$  with  $i_1 < i_2 < \dots < i_k$ , a *spiked cycle*  $C^*$  with  $k$  *spikes* at  $\{i_1, \dots, i_k\}$  arises from the cycle  $C : u_1 u_2 \dots u_\ell u_1$  of length  $\ell$  by attaching a new endvertex  $v_{i_j}$  to  $u_{i_j}$  for every  $j \in [k]$ . The spiked cycle  $C^*$  is *good* if either  $k = 1$  and  $\ell \equiv 1 \pmod{3}$  or  $k \geq 2$ ,

- $i_{j+1} - i_j \equiv 2 \pmod{3}$  for every  $j \in [k-1]$ , and
- $\ell + i_1 - i_k \equiv 1 \pmod{3}$ ,

that is, the  $k$  paths in  $C^*$  between vertices of degree 3 whose internal vertices have degree 2, have lengths  $2, \dots, 2$ , and 1 modulo 3. The spiked cycle  $C^*$  is *very good* if it is good and

- $\ell \not\equiv 1 \pmod{3}$ ,

that is, in particular,  $k \geq 2$ . See Figure 2 for an illustration.




Figure 2: A very good spiked cycle with  $\ell = 15$  and  $k = 5$  spikes at  $\{i_1, \dots, i_k\} = \{2, 4, 9, 11, 13\}$ . Note that removing  $v_4$  results in a spiked cycle that is not good.

Let  $\mathcal{C}$  be the set of all graphs that arise from the graphs in

$$\mathcal{C}_0 = \{P_3\} \cup \{C_\ell : \ell \in \mathbb{N}, \ell \geq 3, \text{ and } \ell \not\equiv 0 \pmod{3}\} \cup \{C^* : C^* \text{ is a very good spiked cycle}\}$$

by repeated applications of the three operation  $(O_1)$ ,  $(O_2)$ , and  $(O_3)$ , as well as the fourth operation  $(O_4)$  of forming the disjoint union of some graph  $G'$  in  $\mathcal{C}$  with a very good spiked cycle  $C^*$ , and adding a bridge between  $V(G')$  and  $V(C^*)$ .

**Lemma 1.** *All graphs in  $\mathcal{C}$  satisfy (2) with equality. Furthermore, for every vertex  $u$  of every graph  $G$  in  $\mathcal{C}$ , the graph  $G$  has a maximum dissociation set not containing  $u$ .*

As our final result, we show that  $\mathcal{C}$  contains all connected cycle-disjoint extremal graphs for Theorem 1. Figure 3 shows two extremal graphs that are not cycle-disjoint.

**Theorem 3.** *A connected cycle-disjoint graph satisfies (2) with equality if and only if it belongs to  $\mathcal{C}$ .*




Figure 3: Two graphs  $G$  that satisfy (2) with equality. Note that the removal of the vertex  $u$ , which lies on two cycles, yields  $d_G(u) - 2$  components.

Within our results, the value  $c_1$  can be replaced by the maximum number of pairwise vertex-disjoint cycles of length 1 modulo 3. It remains to elucidate the structure of all extremal graphs for Theorem 1.

All proofs are provided in the arxiv version of this paper. [2]

## References

- [1] C. Barefoot, Block-cutvertex trees and block-cutvertex partitions, *Discrete Mathematics* 256 (2002) 35-54.
- [2] F. Bock, J. Pardey, L. Penso, and D. Rautenbach, A bound on the dissociation number, *arXiv:2202.09190*
- [3] R. Boliac, K. Cameron, and V.V. Lozin, On computing the dissociation number and the induced matching number of bipartite graphs, *Ars Combinatoria* 72 (2004) 241-253.
- [4] B. Brešar, F. Kardoš, J. Katrenič, and G. Semanišin, Minimum  $k$ -path vertex cover, *Discrete Applied Mathematics* 159 (2011) 1189-1195.
- [5] B. Brešar, M. Jakovac, J. Katrenič, G. Semanišin, and A. Taranenko, On the vertex  $k$ -path cover, *Discrete Applied Mathematics* 161 (2013) 1943-1949.
- [6] F. Göring, J. Harant, D. Rautenbach, and I. Schiermeyer, On  $F$ -independence in graphs, *Discussiones Mathematicae. Graph Theory* 29 (2009) 377-383.
- [7] S. Hosseini and S. Butenko, An improved approximation for maximum  $k$ -dependent set on bipartite graphs, *Discrete Applied Mathematics* 307 (2022) 95-101.
- [8] F. Kardoš, J. Katrenič, and I. Schiermeyer, On computing the minimum 3-path vertex cover and dissociation number of graphs, *Theoretical Computer Science* 412 (2011) 7009-7017.
- [9] Y. Orlovich, A. Dolgui, G. Finke, V. Gordon, and F. Werner, The complexity of dissociation set problems in graphs, *Discrete Applied Mathematics* 159 (2011) 1352-1366.
- [10] E. Speckenmeyer, On feedback vertex sets and nonseparating independent sets in cubic graphs, *Journal of Graph Theory* 12(1988) 405-412.
- [11] D. Tsur, Parameterized algorithm for 3-path vertex cover, *Theoretical Computer Science* 783 (2019) 1-8.
- [12] M. Yannakakis, Node-deletion problems on bipartite graphs, *SIAM Journal on Computing* 10 (1981) 310-327.
- [13] S. Ueno, Y. Kajitani, and S. Gotoh, On the nonseparating independent set problem and feedback set problem for graphs with no vertex degree exceeding three, *Discrete Mathematics* 72 (1988) 355-360.