

A Polynomial-time Algorithm for Outerplanar Diameter Improvement

Nathann Cohen¹, Daniel Gonçalves², Eunjung Kim³, Christophe Paul²,
Ignasi Sau², Dimitrios M. Thilikos², and Mathias Weller²

¹ CNRS, LRI, Orsay, France.

nathann.cohen@gmail.com

² CNRS, LIRMM, Montpellier, France.

{FirstName.FamilyName}@lirmm.fr

³ CNRS, LAMSADE, Paris, France.

eunjungkim78@gmail.com

Abstract. The OUTERPLANAR DIAMETER IMPROVEMENT problem asks, given a graph G and an integer D , whether it is possible to add edges to G in a way that the resulting graph is outerplanar and has diameter at most D . We provide a dynamic programming algorithm that solves this problem in polynomial time. OUTERPLANAR DIAMETER IMPROVEMENT demonstrates several structural analogues to the celebrated and challenging PLANAR DIAMETER IMPROVEMENT problem, where the resulting graph should, instead, be planar. The complexity status of this latter problem is open.

Keywords: diameter improvement; outerplanar graphs; completion problems; polynomial-time algorithms; dynamic programming; parameterized complexity.

1 Introduction

In general, a *graph completion problem* asks, given some i it is possible to add edges to it so that the resulting graph satisfies some target property. There are two different ways of defining the optimization measure for such problems. The first, and most common, is the number of edges to be added, while the second is the value of some graph invariant on the resulting graph. Problems of the first type are HAMILTONIAN COMPLETION [16], INTERVAL GRAPH COMPLETION [18], PROPER INTERVAL GRAPH COMPLETION [17,20], CHORDAL GRAPH COMPLETION [20,24], and STRONGLY CHORDAL GRAPH COMPLETION [20], where the property is being Hamiltonian, interval, proper interval, chordal, and strongly chordal, respectively.

We focus our attention to the second category of problems where, for some given parameterized graph property \mathcal{P}_k , the problem asks, given a graph G and an integer k , whether it is possible to add edges to G such that the resulting graph belongs to \mathcal{P}_k . Usually \mathcal{P}_k is a parameterized graph class whose graphs are typically required (for every k) to satisfy some sparsity condition. There are few problems of this type in the bibliography. Such a completion problem is the PLANAR DISJOINT PATHS COMPLETION problem that asks, given a plane graph and a collection of k pairs of terminals, whether it is possible to add edges such that the resulting graph remains plane and contains k vertex-disjoint paths between each pair of terminals. While this problem is trivially NP-complete, it has been studied from the point of view of parameterized complexity [1]. In particular, when all edges should be added in the same face, it can be solved in $f(k) \cdot n^2$ steps [1], i.e., it is fixed parameter tractable (FPT in short).

Perhaps the most challenging problem of the second category is the PLANAR DIAMETER IMPROVEMENT problem (PDI in short), which was first mentioned by Dejter and Felows [9] (and made an explicit open problem in [12]). Here we are given a planar (or plane) graph and we ask for the minimum integer D such that it is possible to add edges so that the resulting graph is a planar graph with diameter at most D (according to the general formalism, for each D , the parameterized property \mathcal{P}_D contains all planar graphs with diameter at most D). The computational complexity of PLANAR DIAMETER IMPROVEMENT is open, as it is not

even known whether it is an NP-complete problem, even in the case where the embedding is part of the input. Interestingly, PLANAR DIAMETER IMPROVEMENT is known to be FPT: it is easy to verify that, for every D , its YES-instances are closed under taking minors¹ which, according to the meta-algorithmic consequence of the Graph Minors series of Robertson and Seymour [21,22], implies that PLANAR DIAMETER IMPROVEMENT is FPT. Unfortunately, this implication only proves the *existence* of such an algorithm for each D , while it does not give any way to construct it. A uniform FPT algorithm for this problem remains as one of the most intriguing open problems in parameterized algorithm design. To our knowledge, when it comes to explicit algorithms, it is not even clear how to get an $O(n^{f(D)})$ -algorithm for this problem (in parameterized complexity terminology, such an algorithm is called an XP-algorithm).

Notice that, in both aforementioned problems of the second type, the planarity of the graphs in \mathcal{P}_D is an important restriction, as it is essential for generating a non-trivial problem; otherwise, one could immediately turn a graph into a clique that trivially belongs to \mathcal{P}_1 . For practical purposes, such problems are relevant where instead of generating few additional links, we mostly care about maintaining the network topology. The algorithmic and graph-theoretic study on diameter improvement problems has focused both on the case of minimizing the number (or weight) of added edges [2,3,6,11,14,19], as well as on the case of minimizing the diameter [3,15]. In contrast, the network topology, such as acyclicity or planarity, as a constraint to be preserved has received little attention in the context of complementing a graph; see for example [14].

In this paper we study the OUTERPLANAR DIAMETER IMPROVEMENT problem, or OPDI in short. An instance of OPDI consists of an outerplanar graph $G = (V, E)$ and a positive integer D , and we are asked to add a set F of missing edges to G so that the resulting graph $G' = (V, E \cup F)$ has diameter at most D , while G' remains outerplanar. Note that we are allowed to add arbitrarily many edges as long as the new graph is outerplanar. Given a graph $G = (V, E)$, we call $G' = (V, E \cup F)$ a *completion* of G .

It appears that the combinatorics of OPDI demonstrate some interesting parallelisms with the notorious PDI problem. We denote by $\mathbf{opdi}(G)$ (resp. $\mathbf{pdi}(G)$) the minimum diameter of an outerplanar (resp. planar) completion of G . It can be easily seen that the treewidth of a graph with bounded $\mathbf{pdi}(G)$ is bounded, while the pathwidth of a graph with bounded $\mathbf{opdi}(G)$ is also bounded. In that sense, the OPDI can be seen as the “linear counterpart” of PDI. We stress that the same “small pathwidth” behavior of OPDI holds even if, instead of outerplanar graphs, we consider any class of graphs with bounded outerplanarity. It also follows that both $\mathbf{pdi}(G)$ and $\mathbf{opdi}(G)$ are trivially 2-approximable. To see this, let G' be a triangulation of a plane (resp. outerplane) embedding of G where, in every face of G , all edges added to it have a common endpoint. It easily follows that, for both graph invariants, the diameter of G' does not exceed twice the optimal value.

Another closely related notion is that of t -spanners. Given a graph $G = (V, E)$, a t -spanner of G is a spanning subgraph $G' = (V, E')$ such that for every pair of vertices, the distance in G' is within a factor t to their distance in G . The measure t for the quality of a spanner is called the *stretch factor*. The problem of identifying a t -spanner, or the t -SPANNER problem, has been extensively studied in geometry as well as in communication network design [4, 5, 7, 13, 23]. A sparse or *planar* t -spanner is of particular importance: in wireless ad hoc networks, certain routing protocols require the network topology to be planar [23]. The notion of t -spanner is, in a sense, dual to OPDI. The former allows losing the established links and the latter allows having additional links while, in both cases, our interest lies in preserving the constraint on the network topology. Note that in both cases, we want to improve (or do not deteriorate a lot) the performance of the network, which is measured by the lags in information transfer such as diameter or stretch factor.

¹ To see this, if a graph G can be completed into an outerplanar graph G' of diameter D , then G' is also a valid completion of any subgraph $H \subseteq G$. Similarly, by merging two adjacent vertices uv in both G and G' , one sees that the diameter is also closed under edge contraction.

Our results. In this work, we show that OUTERPLANAR DIAMETER IMPROVEMENT is polynomial-time solvable. Our algorithm, described in Section 2, is based on dynamic programming and works in full generality, even when the input graph may be disconnected. Also, our algorithm does *not* assume that the input comes with some specific embedding (in the case of an embedded input, the problem becomes considerably easier to solve).

Further research. Our algorithm for OPDI runs in time $O(n^3)$ for connected input graphs, and in time $O(n^7)$ or $O(n^9)$ for disconnected input graphs, depending on whether D is odd or even, which can probably be improved. By the Graph Minors series of Robertson and Seymour [21,22] we know that for each fixed integer D , the set of *minor obstructions* of OPDI is *finite*. We have some preliminary results in this direction, but we managed to obtain a complete list only for small values of D . We believe that our approach might be interesting for generalizations of the OPDI problem, such as the one where we demand that the completed graph has fixed outerplanarity. Settling the computational complexity of PDI remains the main open problem in this area. An explicit FPT-algorithm, or even an XP-algorithm, would also be significant. Again, we have some partial NP-completeness results in this direction, but for problems which are slightly more complicated than PDI, in particular involving edge weights.

2 Description of the algorithm

The aim of this section is to describe a polynomial-time dynamic program that, given an outerplanar graph G and an integer D , decides whether G admits an outerplanar completion with diameter at most D . Note that such an algorithm easily yields a polynomial-time algorithm to compute, given an outerplanar graph G , the smallest integer D such that G admits an outerplanar completion with diameter D .

Before describing the algorithm, we show some properties of outerplanar completions. In particular, Subsection 2.1 handles the case where the input outerplanar graph has cut vertices. Its objective is to prove that we can apply a *reduction rule* to such a graph which is safe for the OPDI problem. In Subsection 2.2 we deal with 2-vertex separators, and in Subsection 2.3 we present a polynomial-time algorithm for *connected* input graphs. Finally, due to space limitations, we present the algorithm for disconnected input graphs in Appendix A.

Some notation. We use standard graph-theoretic notation, see for instance [10]. It is well known that a graph is outerplanar if and only if it excludes K_4 and $K_{2,3}$ as a minor. An outerplanar graph is *triangulated* if all its inner faces (in an outerplanar embedding) are triangles. An outerplanar graph is *maximal* if it is 2-connected and triangulated. Note that, when solving the OPDI problem, we may always assume that the completed graph G' is maximal. For us, a *block* in a graph is either a 2-connected component or an edge. Given a graph G and a subset $S \subseteq V(G)$, we denote by $\partial(S)$ the set of vertices in S that have at least one neighbor in $V(G) \setminus S$.

2.1 Reducing the input graph when there are cut vertices

Given a graph G , let the *eccentricity* of a vertex u be $\text{ecc}(u, G) = \max_{v \in V(G)} \text{dist}_G(u, v)$. Given an outerplanar graph G , a vertex $u \in V(G)$, and an integer D , let us define $\text{ecc}_D^*(u, G)$ as $\min_H \text{ecc}(u, H)$, where the minimum is taken over all the outerplanar completions H of G of diameter at most D . If all the outerplanar completions have diameter more than D , we set this value to ∞ . Unless said otherwise, we assume henceforth that D is a fixed given integer, so we may just write $\text{ecc}^*(u, G)$ instead of $\text{ecc}_D^*(u, G)$. (The value of D will change only in the description of the algorithm at the end of Subsection 2.3, and in that case we will make the notation explicit).

As admitting an outerplanar completion with bounded eccentricity is a minor-closed property, let us observe the following:

Lemma 1. *For any connected outerplanar graph G , any vertex $v \in V(G)$, and any connected subgraph H of G with $v \in V(H)$, we have that $\text{ecc}^*(v, H) \leq \text{ecc}^*(v, G)$.*

Proof: Let G' be an outerplanar completion of G achieving $\text{ecc}^*(v, G)$, that is such that $\text{ecc}(v, G') = \text{ecc}^*(v, G)$. Contracting the edges of G' that have at least one end out of $V(H)$ one obtains an outerplanar completion H' of H (as outerplanar graphs are minor-closed). As contracting an edge does not elongate any shortest path, the diameter of H' is at most the diameter of G' , and $\text{dist}_{H'}(v, u) \leq \text{dist}_{G'}(v, u)$ for any vertex $u \in V(H)$. We thus have that $\text{ecc}^*(v, H) \leq \text{ecc}(v, H') \leq \text{ecc}(v, G') = \text{ecc}^*(v, G)$. \square

Consider a connected graph G with a separating vertex v , and let C_1, \dots, C_t be the vertex sets of the connected components of $G \setminus \{v\}$. For $1 \leq i \leq t$, we call the vertex set $B_i = C_i \cup \{v\}$ a *branch* of G at v . To shorten notations, we abbreviate $B_i \cup \dots \cup B_j =: B_{i\dots j}$, for any $1 \leq i < j \leq t$. Also, when referring to the eccentricity, we simply denote $G[B_i]$ by B_i . Thus, for example, when considering the value $\text{ecc}^*(v, B_{1\dots i})$, it will refer to the minimum eccentricity with respect to v that an outerplanar completion of the graph $G[B_{1\dots i}]$ with diameter at most D can have.

The following lemma, which is crucial in order to obtain a polynomial-time algorithm, implies that if G has a cut vertex v with many branches, it is safe to remove most of them.

Lemma 2. *Consider an outerplanar graph G with a separating vertex v that belongs to at least 7 branches. Denote these branches B_1, \dots, B_t , with $t \geq 7$, in such a way that $\text{ecc}^*(v, B_1) \geq \text{ecc}^*(v, B_2) \geq \dots \geq \text{ecc}^*(v, B_t)$. The graph G has an outerplanar completion with diameter at most D if and only if $\text{ecc}^*(v, B_{1\dots 6}) + \text{ecc}^*(B_7) \leq D$.*

Proof: “ \Leftarrow ” If $\text{ecc}^*(v, B_{1\dots 6}) + \text{ecc}^*(B_7) \leq D$, gluing on v the outerplanar completions of $G[B_{1\dots 6}], G[B_7], \dots, G[B_t]$, respectively achieving $\text{ecc}^*(v, B_{1\dots 6}), \text{ecc}^*(B_7), \dots, \text{ecc}^*(B_t)$, one obtains an outerplanar completion G' of G with diameter at most D . Indeed,

- The graph obtained is outerplanar and contains G .
- Two vertices x, y of $G[B_{1\dots 6}]$ (resp. of $G[B_i]$ for $7 \leq i \leq t$) are at distance at most D from each other, as $\text{ecc}^*(v, B_{1\dots 6}) < \infty$ (resp. as $\text{ecc}^*(v, B_i) < \infty$).
- Any vertex x of $G[B_{1\dots 6}]$ and y of $G[B_i]$, with $7 \leq i \leq t$, are respectively at distance at most $\text{ecc}^*(v, B_{1\dots 6})$ and $\text{ecc}^*(v, B_i) \leq \text{ecc}^*(v, B_7)$ from v . They are thus at distance at most $\text{ecc}^*(v, B_{1\dots 6}) + \text{ecc}^*(v, B_7) \leq D$ from each other.
- Any vertex x of $G[B_i]$ and y of $G[B_j]$, with $7 \leq i < j \leq t$, are respectively at distance at most $\text{ecc}^*(v, B_i) \leq \text{ecc}^*(v, B_1) \leq \text{ecc}^*(v, B_{1\dots 6})$ (By Lemma 1) and $\text{ecc}^*(v, B_j) \leq \text{ecc}^*(v, B_7)$ from v . They are thus at distance at most D from each other.

“ \Rightarrow ” In the following, we consider towards a contradiction an outerplanar graph G admitting an outerplanar completion with diameter at most D , but such that

$$\text{ecc}^*(v, B_{1\dots 6}) + \text{ecc}^*(v, B_7) > D. \quad (1)$$

Among the triangulated outerplanar completions of G with diameter at most D , let G' be one that minimizes the number of edges added among distinct branches of G . Let us say that these edges are *lianas*. Let t' be the number of branches at v in G' , and denote these branches $B'_1, \dots, B'_{t'}$, in such a way that $\text{ecc}^*(v, B'_1) \geq \text{ecc}^*(v, B'_2) \geq \dots \geq \text{ecc}^*(v, B'_{t'})$. Then, since G' has diameter at most D and shortest paths among distinct branches of G' contain v ,

$$\forall_{1 \leq i' < j' \leq t'} \text{ecc}^*(v, B'_{i'}) + \text{ecc}^*(v, B'_{j'}) \leq D \quad (2)$$

is clear. Let $S_{i'} := \{i \mid B_i \subseteq B'_{i'}\}$ for all $1 \leq i' \leq t'$ and note that $\{S_1, \dots, S_{t'}\}$ is a partition of $\{1, \dots, t\}$. The branches $B'_{i'}$ with $|S_{i'}| = 1$ are called *atomic*. Observe that an atomic branch does not contain any liana. In contrast, for a non-atomic branch $B'_{i'}$, any B_i with $i \in S_{i'}$ is incident to at least one liana in G' in order to connect the graph $G'[B'_{i'} \setminus v]$.

Claim 1 Let $B'_{i'}$ be a non-atomic branch and let $S' \subsetneq S_{i'}$. Then, $\text{ecc}^*(v, \bigcup_{i \in S'} B_i) + \text{ecc}^*(v, \bigcup_{i \in S_{i'} \setminus S'} B_i) > D$.

Proof: Let $\mathcal{B} := \bigcup_{i \in S'} B_i$ and $\bar{\mathcal{B}} := B'_{i'} \setminus \mathcal{B}$. If the claim is false, then $\text{ecc}^*(v, \mathcal{B}) + \text{ecc}^*(v, \bar{\mathcal{B}}) \leq D$. Furthermore, for all $j' \neq i'$,

$$\text{ecc}^*(v, \mathcal{B}) + \text{ecc}^*(v, B'_{j'}) \stackrel{\text{Lemma 1}}{\leq} \text{ecc}^*(v, B'_{i'}) + \text{ecc}^*(v, B'_{j'}) \stackrel{(2)}{\leq} D$$

and, likewise, $\text{ecc}^*(v, \bar{\mathcal{B}}) + \text{ecc}^*(v, B'_{j'}) \leq D$. Thus, the result of replacing $G'[B'_{i'}]$ with the disjoint union of an outerplanar completion achieving $\text{ecc}^*(v, \mathcal{B})$ and an outerplanar completion achieving $\text{ecc}^*(v, \bar{\mathcal{B}})$ yields a diameter- D outerplanar completion containing less lianas than G' , contradicting our choice of G' . \square

Claim 2 Let $2 \leq i' \leq t'$. Then, the branch $B'_{i'}$ is atomic.

Proof: Towards a contradiction, assume that $B'_{i'}$ is not atomic and let $i \in S_{i'}$. Then,

$$\begin{aligned} \text{ecc}^*(v, B'_{i'} \setminus (B_i \setminus v)) + \text{ecc}^*(v, B_i) &\stackrel{\text{Lemma 1}}{\leq} 2 \text{ecc}^*(v, B'_{i'}) \\ &\stackrel{i' > 1}{\leq} \text{ecc}^*(v, B'_1) + \text{ecc}^*(v, B'_{i'}) \stackrel{(2)}{\leq} D, \end{aligned}$$

contradicting Claim 1. \square

In the following, we abbreviate $|S_1| =: s$.

Claim 3 $S_1 = \{1, \dots, s\}$.

Proof: First, assume $1 \notin S_1$. Then, by Claim 2, there is some i' with $B'_{i'} = B_1$. Let j' such that $2 \in S_{j'}$. Let G'' denote the result of, for each j' , replacing $G'[B'_{j'}]$ with an outerplanar completion of B_j achieving $\text{ecc}^*(v, B'_{j'})$. The diameter of G'' is

$$\begin{aligned} \max_{i \neq j} \{ \text{ecc}^*(v, B_i) + \text{ecc}^*(v, B_j) \} &= \text{ecc}^*(v, B_1) + \text{ecc}^*(v, B_2) \\ &\stackrel{\text{Lemma 1}}{\leq} \text{ecc}^*(v, B'_{i'}) + \text{ecc}^*(v, B'_{j'}) \stackrel{(2)}{\leq} D, \end{aligned}$$

contradicting Equation (1).

Second, assume that there is some $i \notin S_1$ with $i+1 \in S_1$. By Claim 2, there is some $i' > 1$ with $B'_{i'} = B_i$. Then,

$$\begin{aligned} \text{ecc}^*(v, B'_1 \setminus (B_{i+1} \setminus v)) + \text{ecc}^*(v, B_{i+1}) &\stackrel{\text{Lemma 1}}{\leq} \text{ecc}^*(v, B'_1) + \text{ecc}^*(v, B_{i+1}) \\ &\leq \text{ecc}^*(v, B'_1) + \text{ecc}^*(v, B_i) \\ &\leq \text{ecc}^*(v, B'_1) + \text{ecc}^*(v, B'_{i'}) \stackrel{(2)}{\leq} D, \end{aligned}$$

contradicting Claim 1. \square

Claim 4 Let $1 < i \leq s$. Then, $\text{ecc}^*(v, B_{1\dots i-1}) + \text{ecc}^*(v, B_i) > D$.

Proof: Since by Claim 3, $B_{1\dots i-1}$ and B_i are subsets of B'_1 , this is a direct consequence of Claim 1 with $S' = \{i, \dots, s\}$. \square

Note that, by (2), $\text{ecc}^*(v, B_{1\dots s}) + \text{ecc}^*(v, B_{s+1}) \leq D$. By (1), this implies $s \geq 7$. Thus, a special case of Claim 4 is $\text{ecc}^*(v, B_1) + \text{ecc}^*(v, B_2) > D$, implying $\text{ecc}^*(v, B_1) > D/2$.

Claim 5 *Let $S' \subseteq \{1, \dots, t\}$ and let $\mathcal{B} := \bigcup_{i \in S'} B_i$. Then, there is a vertex in \mathcal{B} that is, in G' , at distance at least $\text{ecc}^*(v, \mathcal{B})$ to any vertex of $V(G) \setminus (\mathcal{B} \setminus v)$.*

Proof: Let H result from G' by contracting all vertices of $V(G) \setminus \mathcal{B}$ onto v . If the claim is false, then for any vertex $u \in \mathcal{B}$ there exists a vertex $w \in V(G) \setminus (\mathcal{B} \setminus v)$ such that $\text{dist}_{G'}(u, w) < \text{ecc}^*(v, \mathcal{B})$, so contracting all vertices of $V(G) \setminus \mathcal{B}$ onto v yields a path in H between u and v of length strictly smaller than $\text{ecc}^*(v, \mathcal{B})$. As this argument holds for any vertex $u \in \mathcal{B}$, it implies that $\text{ecc}(v, H) < \text{ecc}^*(v, \mathcal{B})$. Since H is an outerplanar completion of $G[\mathcal{B}]$, this contradicts the definition of ecc^* . \square

Two sub-branches B_i and B_j of B'_1 are *linked* if G' has a liana linking them, that is, an edge intersecting both $B_i \setminus \{v\}$ and $B_j \setminus \{v\}$.

Claim 6 *Let $1 \leq i < j \leq s$ and let $\text{ecc}^*(v, B_{1\dots i}) + \text{ecc}^*(v, B_j) > D$. Then, B_j is linked to one of B_1, \dots, B_i .*

Proof: By Claim 5, there is a vertex $x \in B_j$ that is, in G' , at distance at least $\text{ecc}^*(v, B_j)$ to any vertex in $B_{1\dots i}$. Likewise, there is a vertex $y \in B_{1\dots i}$ that is, in G' , at distance at least $\text{ecc}^*(v, B_{1\dots i})$ to any vertex in B_j . Let P be any shortest path of G' between x and y . By Claim 5, the maximal subpath of P in $B_j \setminus v$ containing x has length at least $\text{ecc}^*(v, B_j) - 1$ and the maximal subpath of P in $B_{1\dots i} \setminus v$ containing y has length at least $\text{ecc}^*(v, B_{1\dots i}) - 1$. Since these subpaths are vertex disjoint and, by Claim 4, $\text{ecc}^*(v, B_{1\dots i}) + \text{ecc}^*(v, B_{i+1}) > D$, there is only one edge of P out of these subpaths and this edge intersects both $B_{1\dots i}$ and B_j . This edge realizes the desired link. \square

Claim 4 and Claim 6 clearly imply that for any $1 \leq i < s$, B_{i+1} is linked to one of B_1, \dots, B_i . A consequence of the next claim will be that B_{i+1} is linked to exactly one of these branches.

Claim 7 *The graph (with vertex set $1, \dots, s$) induced by the relation linked (ij is an edge iff B_i is linked to B_j) is a path. Furthermore, the subgraph induced by $\{1, \dots, i\}$, for any $1 \leq i \leq s$, is connected.*

Proof: The first statement is a consequence of the following three facts.

1. This graph is connected. Otherwise, v would be separating in $G'[B'_1]$, contradicting the definition of branch.
2. This graph has maximum degree 2. Consider for contradiction that some branch B_i is linked to three branches B_{j_1} , B_{j_2} , and B_{j_3} . As each of $B_i \setminus v$, $B_{j_1} \setminus v$, $B_{j_2} \setminus v$, and $B_{j_3} \setminus v$ induces a connected graph in G' , these four sets together with v induce a $K_{2,3}$ -minor in G' , contradicting its outerplanarity.
3. This graph is not a cycle. Assume otherwise, so in particular it implies that $s \geq 3$. As each $B_i \setminus v$ induces a connected graph in G' , these sets together with v would induce a K_4 -minor in G' , contradicting its outerplanarity.

The second statement is a direct consequence of the fact that for any $1 \leq i < s$, B_{i+1} is linked to one of B_1, \dots, B_i . \square

Hence, for any $1 \leq i \leq s$, the graph $G'[B_{1\dots i} \setminus v]$ is connected.

Claim 8 *For any $1 \leq i \leq s - 3$, $\text{ecc}^*(v, B_{1\dots i+2}) > \text{ecc}^*(v, B_{1\dots i})$.*

Proof: By Lemma 1, it suffices to show $\text{ecc}^*(v, B_{1\dots i+2}) \neq \text{ecc}^*(v, B_{1\dots i})$. Suppose this is false. Then, Claim 4 implies that $\text{ecc}^*(v, B_{1\dots i}) + \text{ecc}^*(v, B_j) > D$ for all $j \in \{i+1, i+2, i+3\}$. Thus, by Claim 6, each of B_{i+1} , B_{i+2} , and B_{i+3} is linked to one of B_1, \dots, B_i . As each of $B_{1\dots i} \setminus v$, $B_{i+1} \setminus v$, $B_{i+2} \setminus v$, and $B_{i+3} \setminus v$ induces a connected graph in G' , these sets together with vertex v induce a $K_{2,3}$ -minor, contradicting the outerplanarity of G' . \square

In the following let q be any integer such that $4 \leq q \leq s$ and B_q is not linked to B_1 . Note by Claim 8 that $\text{ecc}^*(v, B_{1\dots q-1}) > \text{ecc}^*(v, B_1) > D/2$. Let $p < q$ be such that B_p and B_q are linked. By Claim 7, p is unique. In the following we study the structure of G' between these two branches.

As there is a liana between B_p and B_q and G' is triangulated, there is a path ($v = x_0, x_1, x_2, \dots, x_r$), for some $r \geq 1$, of vertices of B_p (resp. a path ($v = y_0, y_1, y_2, \dots, y_{r'}$), for some $r' \geq 1$, of vertices of B_q) such that a vertex belongs to this path if and only if it is adjacent to a vertex of $B_q \setminus \{v\}$ (resp. $B_p \setminus \{v\}$) in G' . As G' is triangulated, it contains the edge $\{x_1, y_1\}$, and there are $r + r' - 1$ lianas between B_p and B_q .

As for any $0 < i < r$ there is a path from x_{i-1} to x_{i+1} whose inner vertices are y_j 's, note that in the graph $G'[B_{1\dots q-1}]$, the blocks containing the edges $x_{i-1}x_i$ and $x_i x_{i+1}$ are distinct. Indeed, otherwise the vertices x_{i-1}, x_i, x_{i+1} together with a path between x_{i-1} and x_{i+1} in $G'[B_{1\dots q-1}]$, plus the vertex resulting from the contraction of B_q would induce a K_4 -minor, contradicting the outerplanarity of G' . Let X be the block of $G'[B_{1\dots q-1}]$ containing the edge $v x_1$. Note that every branch B_i with $i < q$ and $i \neq p$ belongs to X . Similarly the blocks containing the edges $y_i y_{i+1}$ are all distinct.

Claim 9 *In G' , there is no vertex of $B_{1\dots q-1}$ at distance more than $\text{ecc}^*(v, B_{1\dots q-1})$ to any vertex of B_q .*

Proof: Suppose there exists such a vertex x . As by Claim 5 some vertex y of B_q is at distance $\text{ecc}^*(v, B_q)$ to any vertex of $B_{1\dots q-1}$, the two vertices x, y would be at least $\text{ecc}^*(v, B_{1\dots q-1}) + \text{ecc}^*(v, B_q)$ apart. By Claim 4 this distance is more than D , a contradiction. \square

Claim 10 *Let $x \in B_{1\dots q-1}$ be, in G' , at distance at least $\text{ecc}^*(v, B_{1\dots q-1})$ to any vertex of B_q (including v) — x exists by Claim 5. Then, $x \in X$.*

Proof: Towards a contradiction, assume $x \notin X$. Then, the distance of x and x_1 in G' is at least $\text{ecc}^*(v, B_{1\dots q-1}) - 1$. By Claim 5, there is a vertex $y \in B_1$ (thus $y \in X$) that is, in G' , at distance at least $\text{ecc}^*(v, B_1)$ to both v and x_1 (as $x_1 \in B_p \neq B_1$). As a shortest path between x and y in G' necessarily contains v or x_1 , these vertices are at distance at least $\text{ecc}^*(v, B_{1\dots q-1}) - 1 + \text{ecc}^*(v, B_1)$ to each other. This distance is more than $2\text{ecc}^*(v, B_1) > D$, a contradiction. \square

Claim 11 $\text{ecc}^*(v, B_{1\dots q}) = \text{ecc}^*(v, B_{1\dots q-1})$.

Proof: By Lemma 1, it suffices to show $\text{ecc}^*(v, B_{1\dots q}) \leq \text{ecc}^*(v, B_{1\dots q-1})$. To this end, let H be the outerplanar completion of $B_{1\dots q}$ obtained from G' by contracting every branch B_i , with $i > q$, onto v . Since H is a minor of G' , H is an outerplanar completion of $B_{1\dots q}$ with diameter at most D . We show $\text{ecc}(v, H) \leq \text{ecc}^*(v, B_{1\dots q-1})$.

Let x be a vertex that is, in G' and in H , at distance $\text{ecc}^*(v, B_{1\dots q-1})$ to v , and at distance at least $\text{ecc}^*(v, B_{1\dots q-1}) - 1$ to x_1 . By Claim 10, $x \in X$. As H has diameter at most D , any vertex out of X is either at distance at most $D - \text{ecc}^*(v, B_{1\dots q-1})$ to v , or at distance at most $D + 1 - \text{ecc}^*(v, B_{1\dots q-1})$ to x_1 (and thus at distance at most $D + 2 - \text{ecc}^*(v, B_{1\dots q-1})$ to v). As $\text{ecc}^*(v, B_{1\dots q-1}) > D/2 + 1$, any vertex out of X is thus at distance at most $\text{ecc}^*(v, B_{1\dots q-1})$ from v . Finally Claim 9 implies that the vertices of X are also at distance at most $\text{ecc}^*(v, B_{1\dots q-1})$ from v in H . \square

We now claim that there exist two consecutive such values q between 4 and $s \geq 7$. Indeed, note first that B_1 is linked to at most two other branches, as otherwise these branches together with v and B_1 would induce a $K_{2,3}$ -minor. Note also that by Claim 4 and Claim 6, B_2 is linked

to B_1 , so it follows that B_1 is linked to at most one branch B_j with $j \geq 3$. Therefore, for $4 \leq q \leq 7$, there are at least two consecutive values of q such that B_q is not linked to B_1 , as we wanted to prove.

Once we have two consecutive such values q between 4 and $s \geq 7$, we have by Claim 11 that $\text{ecc}^*(v, B_{1\dots i}) = \text{ecc}^*(v, B_{1\dots i+2})$, for some $i \leq s-2$, contradicting Claim 8. This concludes the proof of the lemma. \square

Our algorithm will compute the eccentricity of a given “root” vertex in a diameter- D outerplanar completion G' of G in which this eccentricity is minimal, that is, $\text{ecc}^*(r, G)$. Then, however, the branch containing the root (B_0 in Algorithm 1, Subsection 2.3) should not be removed. Therefore, although Lemma 2 already implies that G has an outerplanar completion with diameter at most D if and only if $G[B_{1\dots 7}]$ does, we instead use the following corollary to identify removable branches.

Corollary 1. *Let G be an outerplanar graph with a separating vertex v that belongs to at least 8 branches. Denote these branches B_1, \dots, B_t , with $t \geq 8$, in such a way that $\text{ecc}^*(v, B_1) \geq \text{ecc}^*(v, B_2) \geq \dots \geq \text{ecc}^*(v, B_t)$. For each $8 \leq i \leq t$, the graph $G_i = \bigcup_{j \in \{1, \dots, 7, i\}} B_j$ has an outerplanar completion with diameter at most D if and only if G does.*

Proof: Recall that the property of having an outerplanar completion with bounded diameter is minor closed. Thus G_i being a minor of G , we have that if G admits an outerplanar completion with diameter at most D so does G_i .

On the other hand, if G_i admits an outerplanar completion with diameter at most D , by Lemma 2 applied to G_i we have that $\text{ecc}^*(v, B_{1\dots 6}) + \text{ecc}^*(v, B_7) \leq D$. Thus gluing on v the outerplanar completions of $G[B_{1\dots 6}], G[B_7], \dots, G[B_t]$, respectively achieving $\text{ecc}^*(v, B_{1\dots 6}), \text{ecc}^*(v, B_7), \dots, \text{ecc}^*(v, B_t)$, one obtains an outerplanar completion of G with diameter at most D . \square

2.2 Dealing with 2-vertex separators

In this subsection, we extend the definition of eccentricity to the pairs (u, v) such that $uv \in E(G)$. Namely, $\text{ecc}(u, v, G)$ is defined as the set of pairs obtained by taking the maximal elements of the set $\{(\text{dist}_G(u, w), \text{dist}_G(v, w)) \mid w \in V(G)\}$. The pairs are ordered such that $(d_1, d_2) \leq (d'_1, d'_2)$ if and only if $d_1 \leq d'_1$ and $d_2 \leq d'_2$. As u and v are adjacent, note that $\text{dist}_G(u, w)$ and $\text{dist}_G(v, w)$ differ by at most one. Hence, $\text{ecc}(u, v, G)$ equal to one of $\{(d, d)\}, \{(d, d+1)\}, \{(d+1, d)\},$ and $\{(d, d+1), (d+1, d)\}$, for some positive integer d .

Claim 12 *Consider a connected graph $G = (X, E(G))$ with a triangle uvw and two sets $X_u, X_v \subseteq X$ such that $X_u \cup X_v = X$, $X_u \cap X_v = \{w\}$, $\partial(X_u) \subseteq \{u, w\}$, and $\partial(X_v) \subseteq \{v, w\}$. Then $\text{ecc}(u, v, G)$ equals the maximal elements of the set*

$$\begin{aligned} & \{(d_u, \min\{d_u + 1, d_w + 1\}) \mid (d_u, d_w) \in \text{ecc}(u, w, X_u)\} \cup \\ & \{(\min\{d_w + 1, d_v + 1\}, d_v) \mid (d_w, d_v) \in \text{ecc}(w, v, X_v)\}. \end{aligned}$$

Proof: Clear from the fact that a shortest path from $X_u \setminus \{u\}$ to u does not go through $X_v \setminus \{w\}$ (as it should go through $w \in N(u)$), from the fact that a shortest path from X_u to v goes through $\{u, w\} \subseteq N(v)$, and from the fact that any subpath of a shortest path is a shortest path (for some pair of vertices). \square

Given a connected outerplanar graph G , for any two vertices $u, v \in V(G)$ and any vertex set $X \subseteq V(G)$ with $u, v \in X$ such that $\partial(X) \subseteq \{u, v\}$, let us define $\text{ecc}_D^*(u, v, X)$ as the minimal elements of the set

$$\left\{ \text{ecc}(u, v, H) \left| \begin{array}{l} H \text{ is an outerplanar completion of } G[X] \text{ of} \\ \text{diameter at most } D \text{ such that } uv \in E(H) \\ \text{and such that } uv \text{ lies on the outer face.} \end{array} \right. \right\}$$

If this set is empty, we set $\text{ecc}_D^*(u, v, X)$ to (∞, ∞) . Similarly to Subsection 2.1, we may drop the subindex D from the notation $\text{ecc}_D^*(u, v, X)$. Here, $\text{ecc}(u, v, H) \leq \text{ecc}(u, v, H')$ if and only if for any $(d_1, d_2) \in \text{ecc}(u, v, H)$ there exists a $(d'_1, d'_2) \in \text{ecc}(u, v, H')$ such that $(d_1, d_2) \leq (d'_1, d'_2)$. According to the possible forms of $\text{ecc}(u, v, H)$, we have that $\text{ecc}^*(u, v, X)$ is of one of the following five forms:

- $\{(d, d)\}$,
- $\{(d, d+1)\}$,
- $\{(d+1, d)\}$,
- $\{(d, d+1), (d+1, d)\}$, or
- $\{(d, d+1)\}, \{(d+1, d)\}$,

for some positive integer d .

Considering $\text{ecc}^*(u, X)$ for some u and X , note that u has at least one incident edge uv on the outer face in the outerplanar completion achieving $\text{ecc}^*(u, X)$. Thus, we can observe the following.

Observation 1 $\text{ecc}^*(u, X) = \min_{v \in X} \min_{S \in \text{ecc}^*(u, v, X)} \max_{(d_u, d_v) \in S} d_u$.

2.3 The algorithm for connected outerplanar graphs

We now proceed to describe a polynomial-time algorithm that solves OUTERPLANAR DIAMETER IMPROVEMENT when the input outerplanar graph is assumed to be connected. In Appendix A we will deal with the disconnected case. Before proceeding to the formal description of the algorithm, let us provide a high-level sketch.

Algorithm 1 described below receives a *connected* outerplanar graph G , an arbitrary non-separating vertex r of G (called the *root*), and a positive integer D . In order to decide whether G admits an outerplanar completion of diameter at most D , we will compute in polynomial time the value of $\text{ecc}_D^*(r, G)$, which, by definition, is distinct from ∞ if and only if G admits an outerplanar completion of diameter at most D .

In order to compute $\text{ecc}_D^*(r, G)$, the algorithm proceeds as follows. In the first step (lines 1-9), we consider an arbitrary block B of G containing r (line 1), and in order to reduce the input graph G , we consider all vertex separators v of G in B . For each such separator v , we order its corresponding branches according to their eccentricity w.r.t. v (line 8), and by Corollary 1 it is safe to keep just a constant number of them, namely 8 (line 9). For computing the eccentricity of the branches not containing the root (lines 5-7), the algorithm calls itself recursively, by considering the branch as input graph, and vertex v as the new root.

In the second step of the algorithm (lines 10-17), we try all 2-vertex separators u, v in the eventual completed graph G' (note that G' cannot be 3-connected, as otherwise it would contain a $K_{2,3}$ -minor), together with a set X consisting of a subset of the connected components of $G' \setminus \{u, v\}$, not containing the root r . For each such triple (u, v, X) , our objective is to compute the value of $\text{ecc}_D^*(u, v, X)$. For doing so, after initializing its value (lines 11-12), we consider all possible triples w, X_u, X_v chosen as in Claim 12 (line 13), for which we already know the values of $\text{ecc}_D^*(u, w, X_u)$ and $\text{ecc}_D^*(w, v, X_v)$, since the sets X are processed by increasing size. Among all choices of one element in $\text{ecc}_D^*(u, w, X_u)$ and another in $\text{ecc}_D^*(w, v, X_v)$ (line 14), only those whose corresponding completion achieves diameter at most D are considered for updating the value of $\text{ecc}_D^*(u, v, X)$ (line 15). For updating $\text{ecc}_D^*(u, v, X)$ (line 17), we first compute $\text{ecc}_D(u, v, X)$ using Claim 12 (line 16).

Finally, once we have computed all values of $\text{ecc}_D^*(u, v, X)$, we can easily compute the value of $\text{ecc}_D^*(u, X)$ by using Observation 1 (line 18). We can now provide a formal description of the algorithm.

Algorithm 1: OPDI-Connected

Input : A connected outerplanar graph G , a root $r \in V(G)$ such that $G \setminus \{r\}$ is connected, and a positive integer D .

Output: $\text{ecc}_D^*(r, G)$.

// all over the recursive calls of the algorithm, G is a global variable, which gets updated whenever some vertices are removed in line 9

1 Let B be a block of G containing r .

// we consider all 1-separators of B and we reduce G

2 **foreach** separating vertex $v \in V(B)$ **do**

3 Let C_0, \dots, C_t be the connected components of $G \setminus \{v\}$, where $r \in C_0$.

4 Let $B_0 \leftarrow C_0$. // the branch containing the root

5 **for** $i \leftarrow 1$ **to** t **do**

6 Let $B_i \leftarrow G[C_i \cup \{v\}]$. // the branches around v

7 $\text{ecc}_i \leftarrow \text{OPDI-Connected}(B_i, v, D)$. // recursive call to compute $\text{ecc}_D^*(v, B_i)$

8 Reorder the B_i 's so that $\text{ecc}_1 \geq \text{ecc}_2 \geq \dots \geq \text{ecc}_t$.

9 Remove B_8, \dots, B_t from G . // by Corollary 1

// we now guess all 2-separators u, v in the target completion G' , together with a subset X of the connected components of $G' \setminus \{u, v\}$

10 **foreach** triple (u, v, X) such that $r \notin X$ and $\partial(X) \subseteq \{u, v\}$ **do**

 // by increasing size of X , and only if the triple (u, v, X) has not already been considered before in a previous iteration

11 $\text{Tab}_{\text{ECC}}(u, v, X) \leftarrow (\infty, \infty)$. // it corresponds to $\text{ecc}_D^*(u, v, X)$

12 **if** $X = \{u, v\}$ **then** $\text{Tab}_{\text{ECC}}(u, v, X) \leftarrow \{(0, 1), (1, 0)\}$.

13 **else** **foreach** w, X_u, X_v chosen as in Claim 12 **do**

 // eccentricities of smaller subgraphs have been already computed

14 **foreach** $S_u \in \text{Tab}_{\text{ECC}}(u, w, X_u)$ and $S_v \in \text{Tab}_{\text{ECC}}(w, v, X_v)$ **do**

15 **if** for all $(d_u, d_w^u) \in S_u$ and $(d_v, d_w^v) \in S_v$, we have $(d_w^u + d_w^v \leq D)$ or $(d_u + 1 + d_v \leq D)$ **then**

 // if the diameter of the considered completion of X does not exceed D , we compute $\text{ecc}_D(u, v, X)$ using Claim 12

16 $\text{Ecc} \leftarrow \max\{ \{(d_u, \min\{d_u + 1, d_w + 1\}) \mid (d_u, d_w) \in S_u\} \cup \{(\min\{d_v + 1, d_w + 1\}, d_v) \mid (d_w, d_v) \in S_v\} \}$.

 // we now update $\text{ecc}_D^*(u, v, X)$

17 $\text{Tab}_{\text{ECC}}(u, v, X) \leftarrow \min\{\text{Tab}_{\text{ECC}}(u, v, X) \cup \text{Ecc}\}$.

// finally, we compute $\text{ecc}_D^*(r, G)$ using Observation 1

18 **return** $\min_{v \in V(G) \setminus \{r\}} \min_{S \in \text{Tab}_{\text{ECC}}(r, v, V(G))} \max_{(d_u, d_v) \in S} d_u$.

The correctness of Algorithm 1 follows from the results proved in Subsections 2.1 and 2.2. It remains to analyze its running time.

Running time analysis of Algorithm 1. Note that at line 6 each B_i is recursively replaced by an equivalent (by Corollary 1) subgraph such that its separating vertices have at most 8 branches attached.

Let us first focus on the second step of the algorithm, that is, on lines 10-17. The algorithm considers in line 10 at most $O(n^2)$ pairs $\{u, v\}$. As each of u and v has at most 7 attached branches avoiding the root, and $G \setminus \{u, v\}$ has at most 2 connected components with vertices adjacent to both u and v (as otherwise G would contain a $K_{2,3}$ -minor), there are at most $2^7 \cdot 2^7 \cdot 2^2 = 2^{16}$ possible choices for assigning these branches or components to X or not. In line 13, the algorithm considers $O(n)$ vertices w . Similarly, as w belongs to at most 7 branches not containing u nor v , there are at most 2^7 choices for assigning these branches to X_u or

X_v . In lines 14-17, the algorithm uses values that have been already computed in previous iterations, as the sets X are considered by increasing order. Note that each of $\text{ecc}_D^*(u, w, X_u)$ and $\text{ecc}_D^*(w, v, X_v)$ contains at most 2 elements, so at most 4 choices are considered in line 14. Again, at most 4 choices are considered in line 15. Therefore, lines 14-17 are executed in constant time.

As for the first step of the algorithm (lines 1-9), the algorithm calls itself recursively. The number of recursive calls is bounded by the number of blocks of G , as by construction of the algorithm each block is assigned a single root. Therefore, the number of recursive calls is $O(n)$. Once the algorithm calls itself and the corresponding branch has no separating vertex other than the root, the algorithm enters in lines 10-17, whose time complexity has already been accounted above. (Note that each triple (u, v, X) is considered only once, and the value of $\text{ecc}_D^*(u, v, X)$ is stored in the tables.)

Finally, in line 18, the algorithm considers $O(n)$ vertices, and for each of them it chooses among constantly many numbers. Summarizing, we have that the algorithm has overall complexity $O(n^3)$.

It is worth mentioning that Algorithm 1 can also compute the actual completion achieving diameter at most D , if any, within the same time bound. Indeed, it suffices to keep track of which edges have been added to G when considering the guessed triangles uvw (recall that we may assume that the completed graph is triangulated).

Theorem 1. *Algorithm 1 solves OUTERPLANAR DIAMETER IMPROVEMENT for connected input graphs in time $O(n^3)$.*

Note that we can compute $\text{opdi}(G)$ by calling Algorithm 1 with an arbitrary root $r \in V(G)$ for increasing values of D .

Corollary 2. *Let G be a connected outerplanar graph. Then, $\text{opdi}(G)$ can be computed in time $O(n^4)$.*

References

1. I. Adler, S. G. Kolliopoulos, and D. M. Thilikos. Planar disjoint-paths completion. In *Proceedings of the 6th International Symposium on Parameterized and Exact Computation (IPEC)*, pages 80–93, 2011.
2. N. Alon, A. Gyarfas, and M. Ruzinko. Decreasing the diameter of bounded degree graphs. *Journal of Graph Theory*, 35(3):161–172, 2000.
3. D. Bilo, L. Guala, and G. Proietti. Improved approximability and non-approximability results for graph diameter decreasing problems. *Theoretical Computer Science*, 417:12–22, 2012.
4. L. Cai and D. G. Corneil. Tree spanners. *SIAM Journal on Discrete Mathematics*, 8(3):359–387, 1995.
5. B. Chandra, G. Das, G. Narasimhan, and J. Soares. New sparseness results on graph spanners. *International Journal of Computational Geometry & Applications*, 5:125–144, 1995.
6. V. Chepoi, B. Estellon, K. Nouioua, and Y. Vaxes. Mixed covering of trees and the augmentation problem with odd diameter constraints. *Algorithmica*, 45(2):209–226, 2006.
7. E. Cohen. Fast algorithms for constructing t -spanners and paths with stretch t . *SIAM Journal on Computing*, 28(1):210–236, 1998.
8. P. Dankemann, D. Erwin, W. Goddard, S. Mukwembi, and H. Swart. A characterisation of eccentric sequences of maximal outerplanar graphs. *The Australasian Journal of Combinatorics*, 58:376pp, 2014.
9. I. J. Dejter and M. Fellows. Improving the diameter of a planar graph. Technical report, Computer Science department at the University of Victoria, May 1993.
10. R. Diestel. *Graph Theory*. Springer-Verlag, Berlin, 3rd edition, 2005.
11. Y. Dodis and S. Khanna. Design networks with bounded pairwise distance. In *Proceedings of the 31st Annual ACM Symposium on Theory of Computing (STOC)*, pages 750–759, 1999.
12. R. G. Downey and M. R. Fellows. *Parameterized Complexity*. Springer-Verlag, 1999.

13. F. F. Dragan, F. V. Fomin, and P. A. Golovach. Spanners in sparse graphs. In *Proceedings of the 35th International Colloquium on Automata, Languages and Programming (ICALP)*, pages 597–608, 2008.
14. P. Erdős, A. Gyárfás, and M. Ruszinkó. How to decrease the diameter of triangle-free graphs. *Combinatorica*, 18(4):493–501, 1998.
15. F. Frati, S. Gaspers, J. Gudmundsson, and L. Mathieson. Augmenting graphs to minimize the diameter. In *Proceedings of the 24th International Symposium on Algorithms and Computation (ISAAC)*, pages 383–393, 2013.
16. M. R. Garey and D. S. Johnson. *Computers and intractability. A guide to the theory of NP-completeness*. W. H. Freeman and Co., 1979.
17. M. C. Golumbic, H. Kaplan, and R. Shamir. On the complexity of DNA physical mapping. *Advances in Applied Mathematics*, 15(3):251–261, 1994.
18. P. Heggenes, C. Paul, J. A. Telle, and Y. Villanger. Interval completion with few edges. In *Proceedings of the 39th Annual ACM Symposium on Theory of Computing (STOC)*, pages 374–381, 2007.
19. T. Ishii. Augmenting outerplanar graphs to meet diameter requirements. *Journal of Graph Theory*, 74(4):392–416, 2013.
20. H. Kaplan, R. Shamir, and R. E. Tarjan. Tractability of parameterized completion problems on chordal, strongly chordal, and proper interval graphs. *SIAM Journal on Computing*, 28(5):1906–1922, 1999.
21. N. Robertson and P. D. Seymour. Graph Minors. XIII. The disjoint paths problem. *Journal of Combinatorial Theory, Series B*, 63(1):65–110, 1995.
22. N. Robertson and P. D. Seymour. Graph minors. XX. Wagner’s conjecture. *Journal of Combinatorial Theory, Series B*, 92(2):325–357, 2004.
23. Y. Wang and X.-Y. Li. Localized construction of bounded degree and planar spanner for wireless ad hoc networks. *MONET*, 11(2):161–175, 2006.
24. M. Yannakakis. Computing the minimum fill-in is NP-complete. *SIAM Journal on Algebraic and Discrete Methods*, 2(1):77–79, 1981.

A The algorithm for disconnected outerplanar graphs

In this subsection we will focus on the case where the input outerplanar graph is disconnected. The *radius* of a graph is defined as the eccentricity of a “central” vertex, that is, the minimum eccentricity of any of its vertices.

Lemma 3 ([8], Theorem 3). *Let G be a maximal outerplanar graph of diameter D and radius r . Then, $r \leq \lfloor D/2 \rfloor + 1$.*

In the following, we denote the minimum radius of a diameter- D outerplanar completion of a graph or connected component G by $r^*(G)$. If G has no diameter- D outerplanar completion, then let $r^*(G) = \infty$.

Definition 1. *Let G be a connected graph and let D be an integer. Let G' be the graph resulting from G by adding an isolated vertex v . Let G^* be a diameter- D outerplanar completion of G' that minimizes the eccentricity of v . Then, G^* is called escalated completion of (G, D) with respect to v and the eccentricity $\text{ecc}(v, G^*)$, denoted by $r^+(G)$, is called escalated eccentricity of (G, D) . Again, if such a G^* does not exist, let $r^+(G) = \infty$.*

We will apply Definition 1 also to connected components of a graph and, if clear from context, we omit D . Note that we can compute $r^+(G)$ by guessing an edge between the isolated vertex v and G and running `OPDI-Connected`, the algorithm for connected graphs. Hence this can be done in $O(n^4)$ time. Also note that $r^*(G) \leq r^+(G) \leq r^*(G) + 1$.

Lemma 4. *Given a graph G with a connected component C such that $r^+(C) < D/2$, then G has a diameter- D outerplanar completion if and only if $G \setminus C$ does.*

Proof: In a diameter- D outerplanar completion of $G \setminus C$ there is a vertex v with eccentricity at most $\lfloor D/2 \rfloor + 1$, by Lemma 3. In this completion, adding the completion of $C + v$ achieving $r^+(C) < D/2$, yields a diameter- D outerplanar completion of G . \square

Observation 2 *Let C be a connected component of G , let G' be an outerplanar completion of G and let C' be a connected component of $G' \setminus C$. Then, there is a vertex $v \in C$ at distance at least $r^+(C)$ to each vertex of C' in G' .*

Proof: Let the result of contracting all vertices in $G' \setminus (C \cup C')$ onto vertices in C and contracting C' onto a single vertex u be G'' . Then, G'' is a subgraph of an outerplanar completion of the result of adding u as isolated vertex to $G'[C]$. By definition, $\text{ecc}(u, G'') \geq r^+(C)$, implying that there is a vertex $v \in C$ at distance at least $r^+(C)$ to u in G'' . Thus, v is at distance at least $r^+(C)$ to each vertex of C' in G' . \square

Observation 2 immediately implies that any cutset separating two connected components C_1 and C_2 of G in G' has distance at least $r^+(C_1)$ and $r^+(C_2)$ to some vertex in C_1 and C_2 , respectively. Thus, these two vertices are at distance at least $r^+(C_1) + r^+(C_2)$ in G' .

Corollary 3. *Let C_1 and C_2 be connected components of G such that $r^+(C_1) + r^+(C_2) > D$ and let G' be a diameter- D outerplanar completion of G . Then, C_1 and C_2 are adjacent in G' , i.e. G' has an edge with an end in C_1 and an end in C_2 .*

Corollary 3 allows us to conclude that all connected components C with $r^+(C) > D/2$ have to be pairwise adjacent in any diameter- D outerplanar completion of G . Thus, there cannot be more than three such components.

Lemma 5. *An outerplanar graph G with more than 3 connected components C such that $r^+(C) > D/2$ has no diameter- D outerplanar completion.*

On the other hand, if G has no connected component C such that $r^+(C) > D/2$, then G necessarily has a diameter- D outerplanar completion.

Proof: The first statement comes from the above comments. The proof of the second statement is similar to the one of Lemma 4. For some component C of G , let v be such that $\text{ecc}(v, C) = r^*(C) \leq r^+(C) \leq D/2$, and complete C in order to achieve this value. Then for the other components C' consider their escalated completion with respect to v . As $r^+(C') \leq D/2$ this graph has diameter at most D . \square

Hence, assume G has $p = 1, 2$ or 3 connected components C such that $r^+(C) > D/2$. By Corollary 3 these p components are pairwise adjacent in the desired completion. Note that with $O(n^{2p-2})$ tries, we can guess $p - 1$ edges connecting all such components into one larger component. Thus, in the following, we assume that there is only one component C with $r^+(C) > D/2$, denoted \mathcal{C}_{\max} .

Lemma 6. *Consider an outerplanar graph G with exactly one connected components \mathcal{C}_{\max} such that $D/2 < r^+(\mathcal{C}_{\max}) < \infty$. If $r^*(\mathcal{C}_{\max}) \leq D/2$, then G necessarily has a diameter- D outerplanar completion.*

Proof: Same proof as Lemma 5. \square

Let us now distinguish two cases according to the parity of D .

Lemma 7. *For odd D , if an outerplanar graph G has at most one component \mathcal{C}_{\max} such that $D/2 < r^+(\mathcal{C}_{\max}) < \infty$, then G has a diameter- D outerplanar completion.*

Proof: Indeed, by Lemma 4 it is sufficient to consider the component \mathcal{C}_{\max} alone. As $r^+(\mathcal{C}_{\max}) < \infty$, \mathcal{C}_{\max} has a diameter- D outerplanar completion, and so does G . \square

The case where D is even is more technical.

Lemma 8. *For even D , Let p and q respectively denote the number of connected components C such that $D/2 < r^+(C) < \infty$ and $r^+(C) = D/2$, of an outerplanar graph G . If $p \geq 2$ and $p + q \geq 5$, then G has no diameter- D outerplanar completion.*

Proof: By Corollary 3, in a diameter- D outerplanar completion G' of G the p components are pairwise adjacent, and any of the q components is adjacent to every of the p ones. For $p = 2$, as $q \geq 3$, this would induce a $K_{2,3}$ -minor in G' , a contradiction. For the other cases, this would induce a K_4 -minor in G' , a contradiction. \square

Lemma 9. *For even D , if an outerplanar graph G has one component, denoted \mathcal{C}_{\max} , such that $D/2 < r^*(\mathcal{C}_{\max}) < \infty$, and at least 4 other components C such that $D/2 \leq r^+(C) < \infty$, then G has no diameter- D outerplanar completion.*

Proof: Let us denote C_1, C_2, C_3 , and C_4 the connected components such that $r^+(C_i) \geq D/2$, distinct from \mathcal{C}_{\max} . Assume for contradiction that G admits a diameter- D outerplanar completion, denoted G' .

Claim 13 *For each C_i, C_j , either C_i and C_j are adjacent in G' , or C_i and C_j have a common neighbor in G' .*

Proof: Assume for contradiction that C_i and C_j are not adjacent and do not have a common neighbor in G' . Let us now construct the graph G'' from G' as follows. For any component C of $G' \setminus (C_i \cup C_j)$ that is not adjacent to both C_i and C_j , contract C onto vertices of C_i or C_j (According to the one C is neighboring). As G'' is obtained from G' by contracting edges, G'' also is a diameter- D outerplanar completion (for some graph containing C_i and C_j). Let $N_i := N_{G''}(C_i)$, let $N_j := N_{G''}(C_j)$, and note that $C_i \cap N_j = \emptyset$, $N_i \cap C_j = \emptyset$, and $N_i \cap N_j = \emptyset$. Then, by Observation 2 (as $G'' \setminus C_i$ and $G'' \setminus C_j$ are connected), there are vertices $v_i \in C_i$ and $v_j \in C_j$ at distance at least $D/2$ to each vertex in N_i and N_j , respectively, in G'' . Since N_i and N_j are at distance at least one, v_i and v_j are at distance at least $D + 1$, contradicting G'' having diameter D . \square

Claim 14 *There is a vertex $u \in \mathcal{C}_{\max}$ that is adjacent in G' to 3 of the components C_1, C_2, C_3 , and C_4 .*

Proof: First, note that there is a vertex u and 3 components, say C_1, C_2, C_3 , with $u \in N_{G'}[C_i]$ for all $1 \leq i \leq 3$, since otherwise, there would be internally vertex-disjoint paths between each two of the four components C_i , implying the existence of a K_4 -minor in G' .

If u is neither in \mathcal{C}_{\max} nor in C_i , for $1 \leq i \leq 3$, then, since all the C_i are adjacent to \mathcal{C}_{\max} (by Corollary 3), G' would have a $K_{2,3}$ -minor on the vertex sets $\{u, \mathcal{C}_{\max}\}$ and $\{C_1, C_2, C_3\}$.

Hence, in the following, we assume that $u \in C_1$. Let z be a neighbor of C_1 in \mathcal{C}_{\max} and, for $i \in \{2, 3\}$ let w_i denote a neighbor of C_4 in $N[C_i]$. We note that $w_2 \neq z$ and $w_3 \neq z$, since otherwise, the claim follows and we are done. Furthermore, $w_2 \neq u$ and $w_3 \neq u$, since otherwise there is a $K_{2,3}$ -minor on the vertex sets $\{u, \mathcal{C}_{\max}\}$ and $\{C_2, C_3, C_4\}$. Let $X := (C_4 \cup \{w_2, w_3\}) \setminus (C_2 \cup C_3)$ and note that X is adjacent to C_2 and C_3 , respectively. Let Y be the connected component of $\mathcal{C}_{\max} \setminus \{w_2, w_3\}$ containing z , and note that Y is adjacent to C_1 and X . Finally, since X, Y, C_1, C_2 , and C_3 are pairwise disjoint, G' has a $K_{2,3}$ -minor on the vertex sets $\{X, C_1\}$ and $\{C_2, C_3, Y\}$. \square

Let v denote a vertex of \mathcal{C}_{\max} that is at distance at least $D/2 + 1$ to u in G' and consider the result $G' \setminus \{u\}$ of removing u from G' . Let C denote the connected component of $G' \setminus \{u\}$ that contains v . Towards a contradiction, assume there is a connected component C_i that is adjacent to u but not to C in G' , then all paths between v and any vertex in C_i contain u . Since G' has diameter D , all vertices in C_i are at distance at most $D/2 - 1$ to u in G' , contradicting $r^+(C_i) \geq D/2$. Thus there is a $K_{2,3}$ -minor in G' on the vertex sets $\{C_1, C_2, C_3\}$ and $\{u, X\}$ where X is the connected component of $G' \setminus (C_1 \cup C_2 \cup C_3 \cup \{u\})$ containing v . This concludes the proof of the lemma. \square

Hence, assume G has $q = 0, 1, 2$ or 3 connected components C such that $r^+(C) = D/2$. By Corollary 3 these q components are adjacent to each of the p components such that $r^+(C) > D/2$. Note that with $O(n^{2q})$ tries, we can guess q edges connecting each of the q components to one of the p component. Then we are left with a connected graph, and we can call **OPDI-Connected**.

The algorithm itself. We now describe a polynomial-time algorithm that solves the OUTERPLANAR DIAMETER IMPROVEMENT problem when the input contains a disconnected outerplanar graph. Algorithm 2 described below receives a (disconnected) outerplanar graph G , and a positive integer D . At the beginning, the algorithm computes $r^+(C)$ and $r^*(C)$ for each connected component C of G . For computing $r^+(C)$ the algorithm adds a vertex v , guessing (with $O(n)$ tries) an edge connecting v to C , and then calls **OPDI-Connected** for this component and root v . For computing $r^*(C)$ the algorithm guesses a root u (with $O(n)$ tries), and then calls **OPDI-Connected** for C and root u .

If $r^*(C) = \infty$ for some component C then, as $r^*(G) \geq r^*(C)$, G has no diameter- D outerplanar completion.

Then, as they could be added in a diameter- D outerplanar completion (by Lemma 4), the algorithm removes the components C with small escalated eccentricity, that is those such that $r^+(C) < D/2$.

Then the algorithm tests if there is no component C such that $r^+(C) > D/2$, or if there is only one component C such that $r^+(C) > D/2$, and if $r^*(C) \leq D/2$. In both cases by Lemma 5 and Lemma 6, G is a positive instance.

Then the algorithm tests if there is more than 3 component C such that $r^+(C) > D/2$. In this cases, by Lemma 5, G is a negative instance. Otherwise, G has $p = 1, 2$ or 3 such connected components, and the algorithm guesses $p - 1$ edges (in time $O(n^{2p-2})$) to connect them (as they should be by Corollary 3). For each such graph we compute `OPDI-Connected` to check that this graph has a diameter- D outerplanar completion.

Then the algorithm proceeds differently according to D 's parity. If D is odd, then G is a positive instance (By Lemma 7). If D is even, if G has (still) more than $5 - p$ connected components (by Lemma 8 and Lemma 9), then G is a negative instance. Then we are left with a graph G with $1 + q$ connected components, and again the algorithm guesses q edges (in time $O(n^{2q})$), connecting G . For each of these graphs the algorithm calls `OPDI-Connected(G, v, D)` (for any v) to check whether this graph admits a diameter- D outerplanar completion.

Finally if none of these “guessed” connected graphs has a diameter- D outerplanar completion, then the algorithm concludes that G is a negative instance.

Algorithm 2: OPDI-Disconnected

Input : A disconnected outerplanar graph $G = (V, E)$ and an integer D .
Output: “TRUE” if and only if G has an outerplanar completion with diameter $\leq D$.

```

1 foreach connected component  $C$  of  $G$  do
2    $r^+(C) \leftarrow \infty$ 
3    $r^*(C) \leftarrow \infty$ 
4   foreach  $u \in V(C)$  do
5      $Ecc \leftarrow \text{OPDI-Connected}(C, u, D)$ 
6      $r^*(C) \leftarrow \min\{r^*(C), Ecc\}$ 
7      $C' \leftarrow C$  with add vertex  $v$  and added edge  $uv$ 
8      $Ecc \leftarrow \text{OPDI-Connected}(C', v, D)$ 
9      $r^+(C) \leftarrow \min\{r^+(C), Ecc\}$ 
10  if  $r^+(C) = \infty$  then return FALSE
11  if  $r^+(C) < D/2$  then Remove  $C$  from  $G$ 
12 if  $r^+(C) \leq D/2$  for every  $C$  then return TRUE
13 if  $r^+(C) \leq D/2$  for every  $C$  except one,  $C_{max}$ , and  $r^*(C_{max}) \leq D/2$  then
14   return TRUE
15 if  $r^+(C) > D/2$  for at least 4 conn. component  $C$  then return FALSE
16 foreach choice of edges interconnecting these  $p = 1, 2,$  or 3 connected components do
17   // choose  $p - 1$  edges
18   Let  $C_{max}$  be this new conn. component
19   if  $\text{OPDI-Connected}(C_{max}, v, D) < \infty$  then
20     if  $D$  is odd then return TRUE if  $G$  has more than  $5 - p$  conn. comp. then
21       //  $C_{max}$  and  $q$  conn. comp. such that  $r^+(C) = D/2$ 
22       return FALSE
23     else
24       foreach choice of  $q$  edges connecting  $G$  do
25         if  $\text{OPDI-Connected}(G, v, D) < \infty$  then return TRUE
26         return FALSE
27 return FALSE

```

Theorem 2. Algorithm 2 solves OUTERPLANAR DIAMETER IMPROVEMENT for disconnected input graphs in polynomial time. For odd D the running time is $O(n^7)$, while it is $O(n^9)$ for even D .

Proof: Indeed, the algorithm runs in time $O(n^7)$ for odd D (at most $O(n^4)$ at line 16, times $O(n^3)$ for the call to `OPDI-Connected` in line 18). The algorithm runs in $O(n^{2p+2q+1})$ time for even D ($O(n^{2p-2})$ in line 16, times $O(n^{2q})$ in line 22, times $O(n^3)$ for the call to `OPDI-Connected` in line 23), where p and q respectively denote the number of connected components C such that $r^+(C) > D/2$ and $r^+(C) = D/2$. As $p + q \leq 4$ we are done. \square